

INFORMATION TO USERS

This material was produced from a microfilm copy of the original document. While the most advanced technological means to photograph and reproduce this document have been used, the quality is heavily dependent upon the quality of the original submitted.

The following explanation of techniques is provided to help you understand markings or patterns which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting thru an image and duplicating adjacent pages to insure you complete continuity.
2. When an image on the film is obliterated with a large round black mark, it is an indication that the photographer suspected that the copy may have moved during exposure and thus cause a blurred image. You will find a good image of the page in the adjacent frame.
3. When a map, drawing or chart, etc., was part of the material being photographed the photographer followed a definite method in "sectioning" the material. It is customary to begin photoing at the upper left hand corner of a large sheet and to continue photoing from left to right in equal sections with a small overlap. If necessary, sectioning is continued again — beginning below the first row and continuing on until complete.
4. The majority of users indicate that the textual content is of greatest value, however, a somewhat higher quality reproduction could be made from "photographs" if essential to the understanding of the dissertation. Silver prints of "photographs" may be ordered at additional charge by writing the Order Department, giving the catalog number, title, author and specific pages you wish reproduced.
5. PLEASE NOTE: Some pages may have indistinct print. Filmed as received.

Xerox University Microfilms

300 North Zeeb Road
Ann Arbor, Michigan 48106

74-20,359

SICKO, Linda May, 1948-
PHYSIOLOGICAL AND CYTOLOGICAL ASPECTS OF
PHOSPHATE METABOLISM IN THE BLUE-GREEN ALGA,
PLECTONEMA BORYANUM.

The City University of New York, Ph.D., 1974
Biology

University Microfilms, A XEROX Company, Ann Arbor, Michigan

© 1974

LINDA MAY SICKO

ALL RIGHTS RESERVED

PHYSIOLOGICAL AND CYTOLOGICAL ASPECTS OF PHOSPHATE METABOLISM

IN THE BLUE-GREEN ALGA, PLECTONEMA BORYANUM

by

LINDA MAY SICKO

**A dissertation submitted to the Graduate Faculty
in Biology in partial fulfillment for the degree
of Doctor of Philosophy, the City University of
New York**

1974

This manuscript has been read and accepted for the Executive Committee in Biology in satisfaction of the dissertation requirement for the degree of Doctor of Philosophy.

April 19, 1974

Date

Thomas E. Jensen
Chairman of Examining Committee
Professor Thomas E. Jensen

April 19, 1974

Date

Louis G. Moriber
Executive Officer

Jack J. Valdovinos
Prof. J. Valdovinos

Lehman College
Institution

J. Golubow
Prof. J. Golubow

Lehman College
Institution

William D. Cohen
Prof. Wm. D. Cohen

Hunter College
Institution

Louis G. Moriber
Prof. L.G. Moriber

Brooklyn College
Institution

Eugene F. Stoermer
Prof. E. Stoermer

University of Michigan
Institution

The City University of New York

ABSTRACT

PHYSIOLOGICAL AND CYTOLOGICAL ASPECTS OF PHOSPHATE METABOLISM
IN THE BLUE-GREEN ALGA, PLECTONEMA BORYANUM

by

LINDA MAY SICKO

Advisor: Professor Thomas E. Jensen

The effects of various external phosphate concentrations on physiological and cytological aspects of Plectonema boryanum have been studied. P. boryanum was found to tolerate a wide range of phosphate concentrations, from 1 to 1000 mg of phosphate per liter. Growth of the alga in these concentrations was characterized by changes in the subcellular distribution of phosphorus-containing compounds and in ultrastructural changes which were monitored by transmission electron microscopy.

Culturing the alga in phosphate-free or phosphate deficient medium led to general reductions of phosphate in all cell fractions examined, with the most dramatic decrease in both short and long chain polyphosphates. Cytologically, the phosphate starvation period was characterized by the development of areas of medium electron density and expansion of intrathylakoidal spaces.

Inoculation of the phosphate-starved algae into a medium con-

taining a known amount of phosphate led to increases in all phosphorus-containing fractions, particularly the polyphosphates. Increases in both short and long chain polyphosphates were greater than an order of magnitude. The satisfaction of the "phosphorus debt" was met essentially within an hour. Examination of the cells revealed that the cells develop polyphosphate bodies in the characteristic areas of medium electron density that develop during phosphate starvation or phosphate limitation. X-ray energy dispersive analysis of the polyphosphate bodies confirmed that they are deposits consisting of two major elements, phosphorus and calcium.

The alga was not able to utilize the condensed phosphate, sodium tripolyphosphate, without prior hydrolysis of this compound to orthophosphate.

ACKNOWLEDGEMENTS

Sincerest appreciation is extended first and foremost to Dr. Thomas E. Jensen, who not only proved to be an excellent advisor, but also gave me encouragement and moral support when it was needed so badly. I have worked with Dr. Jensen for the past seven years both as an undergraduate and graduate student and can only express my deepest gratitude for the many opportunities he provided that enabled me to attain this goal. I would also like to thank Dr. Jack G. Valdovinos and Dr. Julius Golubow for their many suggestions during the course of the experiments and for their reading of the manuscript.

Sincerest thanks are extended to the following professors for reading the dissertation and offering many helpful suggestions for the writing: Dr. Louis G. Moriber, Dr. William D. Cohen, Dr. Eugene F. Storzner, and Dr. Richard E. Crang.

The dissertation would not be a reality if not for the encouragement and assistance of the following people, some of whom are pursuing study of the elusive phosphate question: Mr. Robert P. Ayala, Ms. Ilene Carmenaty, Mr. Thomas Delendick, Mrs. Barbara Doonan, Mr. Paul Mankiewicz, and Mr. Peter Russo. Mr. Delendick was kind enough to critically read the manuscript. Mr. Ayala and Ms. Carmenaty were of great assistance in maintaining the algal

cultures and in the preparation of the photographs.

I would also like to thank the following for their technical assistance: Mr. J. R. Condray of the Monsanto Chemical Co., St. Louis Missouri, for supply samples of condensed phosphates, and Mr. Robert Santorelli, Mr. David Harling, Mr. Peter Newel, and Mr. John Bonnicci of the JEOL Co., Medford, Massachusetts, for providing the necessary advice and equipment for the x-ray energy dispersive analysis.

Special thanks and love are extended to my family, especially my grandmother, Mrs. Mary Flasiński, for their love and encouragement for the past four years.

Last, but certainly not least, my deepest appreciation and love is extended to Earl Glen Goad, my fiancée, who was patient enough to wait, who gave me words of encouragement in times of despair, and who waits no longer.

TABLE OF CONTENTS

	Page
1. Acknowledgments	vi
2. List of Tables and Figures	1
3. Introduction	9
4. Literature Review	11
A. Widespread occurrence of phosphorus in nature	11
B. Biological occurrence of polyphosphate	15
C. Parameters affecting active uptake of phosphate	18
D. Microscopy of polyphosphate bodies	22
E. Aspects of phosphate metabolism	27
F. Phosphagen hypothesis	34
G. Polyphosphate as a phosphorus reserve	36
H. Ecological aspects of phosphorus in natural waters.	37
5. Materials and Methods.	50
A. Growth curves	50
B. Dry weight determinations	51
C. Correlation of dry weight to cell counts	51
D. Phosphate assay and calibration	52
E. Starvation conditions	53
F. "Overplus" or rapid uptake	54
G. Inhibitors and temperature studies	54
H. Total phosphate determinations	55
I. Extraction scheme for polyphosphates	56

	Page
i. Cold TCA extraction	56
ii. Lipid extraction	56
iii. Hot TCA extraction	56
iv. Separation of nucleic acids	58
J. Electron microscopy	58
K. Light microscopy	59
L. X-ray energy dispersive analysis	60
M. Determination of standard deviation and standard error. . .	61
6. Results and Observations	62
A. Growth curves	62
B. Total phosphate present as a function of growth	67
C. Determination of optimum starvation length for rapid uptake	74
D. Determination of uptake parameters	79
E. Physical parameters affecting phosphate uptake rates. . . .	85
i. Temperature	85
ii. Light	85
iii. Effect of phosphate concentration on uptake	88
iv. Effects of inhibitors on uptake	88
F. Normal levels of phosphorus containing compounds in <u>P. boryanum</u>	95
G. Phosphorus distributions during starvation and rapid uptake	100
H. Hydrolysis of condensed phosphates	112
I. Rapid uptake of condensed phosphates	117
J. Cell ultrastructure under normal culture conditions	117

	Page
K. Changes in ultrastructure associated with several phosphate concentrations	120
L. Key to figure legends of electron micrographs	121
M. Cells grown in the absence of phosphate	131
N. Changes in ultrastructure during rapid uptake	136
O. Light microscopy	145
P. X-ray energy dispersive analysis	148
7. Discussion	158
A. Growth curves, phosphate depletion, and minimum cellular phosphate levels	158
B. Physical parameters affecting uptake rates	160
C. Phosphorus distributions in cellular extracts	162
D. Hydrolysis and uptake of condensed phosphates	167
E. Electron microscopy	169
8. Summary	176
9. Appendices	179
10. Literature cited	186

LIST OF TABLES

Table	Page
1. Mutants of <u>Aerobacter aerogenes</u> deficient in polyphosphate metabolism	32
2. Sources of nutrients occurring in natural waters	39
3. Changes in cell size and weight during starvation and rapid uptake	82
4. Summary of phosphate depletion in the medium and increase in total cell phosphorus as a function of external orthophosphate depletion	94
5. Phosphorus content of cell fractions as a function of phosphate concentration in the culture medium	99
6. Phosphorus content of cell fractions during 5 days of phosphate starvation and 4 hours of phosphate uptake	102

LIST OF FIGURES

Figure	Page
1. Structures of several condensed phosphates	12
2. Fractionation scheme for the extraction of phosphorus- containing compounds	57
3. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft.-candles of illumination, 12 hour alternating day/night cycle, 1 mg PO ₄ / liter. . .	64
4. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle, 2 mg PO ₄ / liter. . .	64
5. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle, 5 mg PO ₄ / liter. . .	64
6. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle, 8 mg PO ₄ / liter. . .	64
7. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle. 10 mg PO ₄ / liter. .	66
8. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle. 100 mg PO ₄ / liter .	66

Figure	Page
9. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 25°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle. 1000 mg PO ₄ / liter.	66
10. Growth curve of <u>Oscillatoria tenuis</u> . Same parameters as figure 3.	69
11. Growth curve of <u>O. tenuis</u> . Same parameters as figure 4.	69
12. Growth curve of <u>O. tenuis</u> . Same parameters as figure 5.	69
13. Growth curve of <u>O. tenuis</u> . Same parameters as figure 6.	69
14. Growth curve of <u>O. tenuis</u> . Same parameters as figure 7.	71
15. Growth curve of <u>O. tenuis</u> . Same parameters as figure 8.	71
16. Growth curve of <u>O. tenuis</u> . Same parameters as figure 9.	71
17. Growth curve of <u>P. boryanum</u> and phosphate depletion of culture medium. 37°C, 500 ft-candles of illumination, 12 hour alternating day/night cycle, 10 mg PO ₄ / liter	73
18. Total phosphorus content of <u>P. boryanum</u> as a function of culture age. Cultures maintained at 25°C.	76
19. Total phosphorus content of <u>P. boryanum</u> as a function of culture age. Cultures maintained at 37°C.	78
20. Semi-logarithmic plot of the increase in total cell phosphorus as a function of increasing the length of time in phosphate-free medium (starvation)	81
21. Increase in total cell phosphorus and cold TCA extractable phosphorus during rapid uptake	84
22. Increase in dry weight of the algae and depletion of medium phosphorus during rapid uptake.	84

Figure	Page
23. Semi-logarithmic plot of the increase in total cell phosphorus versus uptake time as a function of temperature	87
24. Increase in total cell phosphorus during rapid uptake as a function of light intensity	90
25. Change in total cell phosphorus and phosphate depletion of the medium as a function of orthophosphate concentration in the medium. 6.58 mg PO_4 / liter.	92
26. Change in total cell phosphorus and phosphate depletion of the medium as a function of orthophosphate concentration in the medium. 10.13 mg PO_4 / liter . . .	92
27. Change in total cell phosphorus and phosphate depletion of the medium as a function of orthophosphate concentration in the medium. 92.51 mg PO_4 / liter . . .	92
28. Effect of metabolic inhibitors on the rapid uptake process. Semi-logarithmic plot of increase in total cell phosphorus versus time	97
29. Semi-logarithmic graphic presentation of the change in total cell phosphorus during starvation and rapid uptake	104
30. Semi-logarithmic graphic presentation of the change in cold TCA extractable phosphorus during starvation and rapid uptake.	107

Figure	Page
31. Semi-logarithmic graphic presentation of the change in phosphorus content of the lipid fraction during starvation and rapid uptake	109
32. Semi-logarithmic graphic presentation of the change in hot TCA extractable phosphorus during starvation and rapid uptake	111
33. Semi-logarithmic graphic presentation of the change in phosphorus content of the residue from the threefold extraction during starvation and rapid uptake	114
34. Hydrolysis of sodium tripolyphosphate in Modified Fitzgerald medium both in the presence and absence of <u>P. boryanum</u> . 4.59 mg PO ₄ / liter as condensed phosphate. .	116
35. Hydrolysis of sodium tripolyphosphate in Modified Fitzgerald medium both in the presence and absence of <u>P. boryanum</u> . 153.7 mg PO ₄ / liter as condensed phosphate .	116
36. Hydrolysis of sodium tripolyphosphate in Modified Fitzgerald medium both in the presence and absence of <u>P. boryanum</u> . 1204 mg PO ₄ / liter as condensed phosphate. .	116
37. Increase in phosphorus - containing fractions during starvation and rapid uptake from 10 mg PO ₄ (as sodium tripolyphosphate) per liter	119
38. Cells of <u>Plectonema boryanum</u> grown for 14 days in 10 mg PO ₄ / liter at 500 ft-candles of illumination. 25°C, and an alternating 12 hour day/night cycle.	123

Figure	Page
39. <u>P. boryanum</u> grown under normal culture conditions as listed above, and showing cell wall detail	123
40. Low power picture of a sectioned pellet grown in medium containing 100 mg PO ₄ / liter. Cell debris is evident . . .	125
41. Cells grown in 1000 mg PO ₄ / liter showing some intrathylakoidal vacuolization.	127
42. Cells grown in 1000 mg PO ₄ / liter, devoid of ribosomes, and apparently dying	127
43. Filament of <u>P. boryanum</u> grown in 1000 mg PO ₄ / liter and containing numerous polyphosphate bodies	127
44. Cells grown in 1 mg PO ₄ / liter showing some expansion of intrathylakoidal spaces	130
45. Cells grown in 1 mg PO ₄ / liter showing a greater degree of expansion of intrathylakoidal spaces	130
46. Cell grown in 1 mg PO ₄ / liter which has unusually large ribosomes	130
47. Cell grown in 1 mg PO ₄ / liter, and unusually elongated . .	130
48. Elongated cell of <u>P. boryanum</u> grown in phosphate-free medium for 2 days	133
49. Portion of a filament of <u>P. boryanum</u> starved of phosphate for 3 days and showing both unusual and unequal cell divisions	133
50-52. Cells of <u>P. boryanum</u> grown in phosphate-free medium for 5 days, and showing large expansion of intrathylakoidal spaces and the development of an electron-lucent area . . .	135

Figure	Page
53-54. Two cell types of <u>P. boryanum</u> from a culture starved of phosphate for 3 days, showing the presence of large lipid-like inclusions	138
55. Cell of <u>P. boryanum</u> at 2 hours of phosphate uptake, with a large electron-lucent area in the center of the cell, and some polyphosphate present in the section	140
56. Uptake cell with a large electron-lucent area and DNA at the periphery of this area	140
57. Uptake cell containing unusually large ribosomes.	142
58. Uptake cells with large electron-lucent areas, and numerous polyphosphate bodies in various stages of development	142
59-62. Cells of <u>P. boryanum</u> at 4 hours of uptake showing large polyphosphate bodies.	144
63. Portion of a filament of <u>P. boryanum</u> at 4 hours of uptake showing electron-lucent areas and remnants of polyphosphate bodies	147
64. Light micrograph of a filament starved of phosphate for 5 days and stained for polyphosphates	147
65. Light micrograph of a filament stained for polyphosphates at 4 hours of phosphate uptake.	147
66. X-ray energy dispersive analysis of cytoplasm of an uptake cell containing no polyphosphate bodies. Glutaraldehyde fixed and Epon embedded.	150

Figure	Page
68. X-ray energy dispersive analysis of a developing polyphosphate body. Glutaraldehyde fixed and Epon embedded	150
69. X-ray energy dispersive analysis of epoxy containing no cells. Osmium fixation.	153
70. X-ray energy dispersive analysis of cytoplasm containing no discernible polyphosphate bodies. Osmium fixed and Epon embedded	153
71. X-ray energy dispersive analysis of a dense polyphosphate body. Osmium fixed and Epon embedded	153
72. X-ray energy dispersive analysis of Durcupan embedding medium. Osmium fixation.	156
73. X-ray energy dispersive analysis of cytoplasm containing no discernible polyphosphate bodies. Osmium fixation and Durcupan embedded	156
74. X-ray energy dispersive analysis of a dense polyphosphate body. Osmium fixation and Durcupan embedded.	156

INTRODUCTION

A paradox exists in the apparent success of blue-green algae in eutrophic waters. Blooms of these species often occur when the supply of nutrients is at its lowest point (Pearsall, 1932). Hutchinson (1973) suggests that a mechanism exists in these organisms which makes them extremely efficient at taking up phosphorus at very low concentrations. This suggestion is supported by evidence from Mackereth (1953), Rigler (1956), Stewart and Alexander (1971), and Lean (1973). The mechanism suggested by Hutchinson is most likely a physiological condition referred to by other workers as "Phosphat-Uberkompensation" (Liss and Langen, 1962), "polyphosphate overplus phenomenon" (Harold, 1964, 1965; Harold and Harold, 1963; Voelz et al, 1966), or "rapid uptake" (Shapiro, 1967) and provides for a restoration of phosphorus supply following phosphate starvation or phosphate limitation. The specific mechanism involved results in the formation of polyphosphates which are long chain, osmotically inert polymers of inorganic phosphate.

The biochemical nature of polyphosphates, their periodicity and role in phosphorus metabolism in the cell, and the significance of their presence in algae common in eutrophic waters require further examination. Difficulties arise when one attempts to assess the extensive literature available on phosphate metabolism. Both environmental and physiological parameters of the uptake phenomenon described in the literature vary. In addition, no

exhaustive study has been conducted on one test organism.

Consideration of these factors led to studies which were designed to determine if the polyphosphate overplus phenomenon exists in a blue-green alga, Plectonema boryanum Gomont. The experiments are an attempt to elucidate the effect of environmental parameters on the physiological and cytological responses of the alga to conditions of both phosphate limitation and excess available phosphate.

LITERATURE REVIEW

Widespread Occurrence of Phosphorus in Nature

Phosphorus can be found in a variety of forms in water, the usual environment of algae. These can be classified as follows:

1. orthophosphate, 2. condensed inorganic phosphates, and 3. organic phosphorylated compounds.

Orthophosphate is the most commonly encountered form of phosphate. Uptake of orthophosphate constitutes the basic process of phosphate metabolism in algae as well as other lower organisms (Kuhl, 1968) and is the simplest form of inorganic phosphate.


Condensed inorganic phosphates may be defined as "pentavalent phosphorus compounds in which various numbers of tetrahedral PO_4 groups are linked together by oxygen bridges" (Thilo, 1962). Condensed phosphates may further be subdivided into three classes (Harold, 1966): a. cyclic condensed phosphates, b. linear condensed phosphates (Figure 1), and c. cross-linked condensed phosphates.

Cyclic condensed phosphates have the general elementary formula $M_n P_n O_{3n}$. These phosphates are commonly referred to as metaphosphates. Treatment with strong alkali converts them to their corresponding linear polymer, and they can be further hydrolyzed to orthophosphate when heated in strong acid (Harold, 1966).


Linear condensed phosphates have the elementary formula

Figure 1*


Structures of several condensed phosphates




Pyrophosphoric acid




Triphosphoric acid



Polyphosphoric acid



Trimetaphosphoric acid



Tetrametaphosphoric acid

*after
F.M. Huennekens and H.R. Whitley, 1960

$M_{n+2}P_nO_{3n+1}$, and are commonly referred to as polyphosphates. They range in size from a chain length of 2 (pyrophosphate) to Kurrol's and Maddrell's salts of chain length of around 10^4 (Harold, 1966). Polyphosphate is stable in alkali but labile to acid. Analysis of polyphosphate is achieved by hydrolyzing in the presence of 1N HCl for 7 minutes at 100°C (LePage, 1949).

Cross-linked condensed phosphate or "ultraphosphates" are the third class of inorganic polymers which are characterized by branching points, or phosphate groups in which three oxygen atoms are shared with neighboring phosphate groups. The biological occurrence of ultraphosphate is not encountered since the branching points may be readily hydrolyzed in water (Harold, 1966).

Organic Phosphorylated Compounds

Phosphorus may be found in many forms in cells other than in an inorganic form. These forms are called organic phosphorylated compounds. Examination of these compounds by a scheme such as proposed by Winder and Denney (1956) reveals that the following fractions may be found in cells: a. phospholipid, b. phosphoprotein, c. deoxyribonucleic acid, d. ribonucleic acid, e. phosphorylated "high-energy" compounds such as ATP, and f. phosphorylated intermediates.

Estimation of these compounds can be complicated by the presence of inorganic polymers which fall into two classes, 1. acid soluble, and 2. acid insoluble. These complications will be discussed in the following section.

Chemical Methods of Analysis

Chemical methods of analysis for polyphosphates depend in all cases upon the final acid hydrolysis of the polymer to orthophosphate. The most widely used methods for the extraction of polyphosphates are modifications of the Schmidt-Thannhauser scheme for extraction of nucleic acids (Schmidt-Thannhauser, 1945). There are two major fractions of polyphosphates derived from these schemes; 1. polyphosphates which are soluble in cold (4°C) 5% trichloroacetic acid (TCA) or perchloric acid, and 2. those which are soluble in hot (70°C) 5% trichloroacetic acid or perchloric acid (Harold, 1962; 1966). Those soluble in cold TCA are referred to as "acid-soluble", and those soluble only in hot TCA are "acid-insoluble". The chain lengths of the two fractions vary, shorter chain length polymers being soluble in cold acid, and longer chain polymers being soluble in hot acid (Juni et al 1947; Wiame, 1949). Separation of any inorganic polymers from nucleic acids can be achieved by the use of charcoal (Harold, 1966). Nucleic acids are absorbed to the charcoal and can later be eluted by alcoholic ammonia.

Methods for the estimation of orthophosphate depend on the formation of a phospho-molybdate complex and its reduction to a highly colored blue compound (Harwood, 1969). The most widely used method for the formation of this colored complex is the stannous chloride method (Amer. Publ. Health Assoc., 1965). Complications arise due to a large salt error, interference by arsenate ions, and instability of the complex formed (Hosakawa and Ohshima, 1973). Another widely used and perhaps more reliable method is that of

Murphy and Riley utilizing ascorbic acid. The salt error is considerably less, interference by other ions is almost negligible, and the colored complex remains stable for 24 hours (Murphy and Riley, 1962).

Biological Occurrence of Polyphosphate

Polyphosphate has been a subject of controversy for at least twenty-five years. In 1944, Jeener and Brachet noted a massive accumulation of a basophilic substance within yeast cells after the addition of phosphate to a previously phosphate-starved suspension. Wiame (1947a,b, 1949) and Schmidt et al (1946) isolated this substance and identified it as polyphosphate. Thus, granules earlier referred to a "metachromatic" or "volutin" granules were actually discovered to be deposits of inorganic polyphosphates.

Polyphosphates have been reported to occur in a variety of organisms from bacteria and blue-green algae to higher plants and animals (Harold, 1966). The occurrence of polyphosphates seems to be related to two distinct nutritional conditions (Smith et al, 1954). The first of these is nutrient imbalance. Exhaustion of an essential nutrient can result in the formation of many types of reserve materials in the cell such as poly β -hydroxybutyric acid, glycogen, polyphosphates, or lipids, which are all osmotically inert (Voelz et al, 1966). Pesch (1924) showed that in Corynebacterium diphtheriae volutin production increased when the amount of growth was limited on glucose-blood agar medium due to insufficient blood content. Duguid et al (1954) found in Klebsiella aerogenes that volutin production

was increased on inadequately buffered sugar-containing agar medium. Smith et al (1954) obtained evidence that volutin formation in Aerobacter aerogenes occurred when cultures were limited by nitrogen or sulfur deficiencies, but not carbon, potassium, or phosphorus. In a similar manner, Spitznagel and Sharp (1959) reported that magnesium deficiency interfered with volutin formation, whereas sulfate deficiency promoted volutin formation in Mycobacterium bovis.

The second nutritional condition that can stimulate the formation of polyphosphate is the restoration of a phosphorus supply following phosphate starvation. This phenomenon has been referred to as "Phosphat-Überkompensation" (Liss and Langen, 1962) or "polyphosphate overplus" (Harold, 1964, 1965; Harold and Harold, 1963; Voelz et al, 1966). The ability of cells synthetically to accumulate phosphorus was first investigated by Ketchum (1939) in the marine diatom Nitzschia closterium. He referred to the deficiency as the phosphorus or phosphate debt, and measured this debt by the amount of phosphate absorbed from the medium by the diatom, or by direct analysis of the cells. The magnitude of the phosphorus debt was directly related to the length of time the cells grew in the light in phosphorus-free medium.

Jeener and Brachet (1944) found that in yeast cells, basophilia associated with volutin granules decreased when the cells were grown on a phosphorus-deficient medium, and increased when the cells were transferred to a medium containing phosphorus. Wiame (1947) demonstrated abundant volutin synthesis when phosphate-starved Saccharomyces

cerevisiae was transferred to a phosphate-rich medium. This phenomenon has also been studied in such bacteria as Caulobacter (Gruha et al, 1954), Klebsiella aerogenes (Duguid, 1948), Aerobacter aerogenes (Smith et al, 1954), and Myxococcus xanthus (Voelz et al, 1966).

Blum (1966) studied several parameters of phosphate uptake by phosphate-starved Euglena gracilis. He reported that many features of this uptake system were similar to those of an active transport system. That is, in Euglena, phosphate uptake requires an energy source, can be limited by dinitrophenol, is saturable by substrate, and can be competitively inhibited by a phosphate analog, arsenate. Similar evidence for an active uptake process is presented by Kylin (1966) in Scenedesmus, by Simonis and Urbach (1963) in Ankistrodesmus braunii, and by Borst-Pauwels and Jager (1969) in Saccharomyces cerevisiae.

Two exceptions have been reported regarding "luxury consumption" of phosphate and involvement of an active uptake process. Butt and Lees (1960) have reported that phosphorus deficient cells of Nitrobacter assimilate orthophosphate from the medium at a slow rate during nitrite oxidation. Whitton (1967), studying phosphate accumulation by Nostoc colonies, found that "softer" Nostoc colonies could accumulate phosphate from the environment by non-active means. This accumulation of phosphate was reduced by pretreatment with chelating agents.

Parameters Affecting Active Uptake of Phosphate

As has been demonstrated many times, the uptake of phosphorus in "luxury consumption" is a process requiring energy. This uptake can be limited by physical and chemical parameters such as light, pH, temperature, concentration gradient, source of available phosphorus, ion effects (Fogg, 1973), and other factors collectively referred to as a "phosphate-sparing" factor (Shapiro, 1968).

Light stimulation of absorption of ^{32}P labelled phosphate has been reported by Talpasayi (1962) for the blue-green alga Anabaena cylindrica. Simonis and Urbach (1963) found that preillumination of cultures of the green alga Ankistrodesmus braunii stimulated subsequent assimilation of phosphate in the dark. Other reports of dark assimilation of phosphate have been made in Chlorella by Kanai, Miyachi, and Miyachi (1963), in Anacystis nidulans by Batterton and Van Baalen (1968), in several algae by Stewart and Alexander (1971), and Overbeck (1962) who demonstrated a dark "overplus" phenomenon in Scenedesmus. Light and/or dark uptake of phosphate has been found to exist in Selenastrum capricornutum (Fitzgerald, 1970) and in Ankistrodesmus braunii (Kanai and Simonis, 1968). Harris and Riley (1956) suggested that a dark uptake of phosphate may allow replenishment of phosphorus to phytoplankton which become phosphorus deficient during the day. Fitzgerald (1970) interprets these findings as a means of nutrient absorption by algae in areas of low light intensity. The algae could then rise to the photic zone of a lake where growth could later take place.

Phosphate uptake rates can be affected markedly by such physical parameters as stream current, concentration of other nutrients, and

the aquatic environment itself (lake water vs. laboratory conditions). Fogg (1973) discusses the influence of motion on uptake of nutrients by planktonic species. Those organisms which are non-motile achieve changes in concentration gradient by sinking or by motion of currents. Motile organisms are able to change their position and can seek out optimal growth conditions; however, they still may be affected by currents. Schumacher and Whitford (1965) found that in a variety of algae, stream currents as low as 1-4 cm/sec increased the rate of phosphate uptake. The concentration of phosphorus in natural waters is usually low, ranging from about 1-20 ug of phosphorus per liter. These low concentrations do not always repress growth. Atkins (1923, 1925) found that the diatom Nitzschia closterium grew well in cultures until it completely utilized all the available phosphate. The growth of the diatom in natural waters appeared to be seasonal; the phosphate concentrations became higher in the winter, accounting for the summer development of phytoplankton. Kuenzler and Ketchum (1962) showed that Phaeodactylum tricorutum was able to take up phosphorus from solutions containing less than 11nM phosphate, and suggested that concentration levels of phosphate this low do not depress growth rate. Pilson and Betzer (1973), studying the phosphate flux across a coral reef, found that the phytoplankton take up phosphorus at a nearly constant rate, independent of light quantity, or the magnitude of photosynthetic activity.

The normal low value of phosphate in water has led several authors to discuss the possibility of a "phosphate-sparing" factor (Shapiro, 1968), that is, a substance, quite possibly organic, which

allows lake water organisms to respond quite differently to varying phosphate concentration. In 1948, Rodhe demonstrated that the diatom Asterionella formosa would grow well in lake water with concentrations of phosphate as low as 0.002 mg/l. This same diatom, under laboratory growth conditions and in a medium consisting of inorganic salts in distilled water, would only grow when the phosphate concentration was 0.20 mg/l. Mackereth (1953) investigated the active uptake of phosphate by this diatom, and found that luxury consumption of phosphate occurred in lake water supplemented with phosphate. No uptake occurred in phosphate solutions (30ug/l) made up in distilled water. Mackereth drew no conclusions as to why the diatoms behaved so differently in lake water and in artificial medium. Using a similar system, Shapiro (1968) postulated that in his test organism, Microcystis aeruginosa, the concentration of inorganic anions was probably the most important parameter affecting uptake.

Ion effects and pH can also markedly affect uptake of ortho-phosphate and distribution into the phosphorus-containing compounds of the cell. Both types of experiments indicate that fixed charges on either the plasmalemma or cell wall can prevent an active ion uptake. Ullrich (1972) found that in synchronized cultures of Ankistrodesmus braunii, ³²P - labelling is strongly dependent upon the pH of the culture medium. In alkaline ranges and in the absence of CO₂, organic phosphates and ATP are labelled most strongly, whereas polyphosphate labelling is highest in the acidic range. Using the same organisms, Ullrich-Eberius (1973) reported that

maximal rates of phosphate uptake occurred between pH 5.5 and 6.5, in agreement with Ullrich. Ullrich-Eberius also found that Na^+ enhanced phosphate uptake 8 to 9 times in the light and in the dark. Belsky et al (1970) also reported a specific Na^+ requirement for phosphate uptake in the marine fungus Dermocystidium sp. Ullrich-Eberius and Simonis (1970) investigated the effect of both sodium and potassium ions on phosphate uptake, by A. braunii. Again, 0.002M sodium chloride increased phosphate uptake whereas uptake was constant over a long period of time in the presence of 0.002M KCl. The authors, considering these ion effects coupled with pH dependency, suggested that Ankistrodesmus metabolically transports H_2PO_4^- , but not HPO_4^{2-} , and that these effects are exerted at the plasmalemma.

Other ion effects have also been reported in the literature. Polyphosphate formation in Saccharomyces mellis is almost completely inhibited by 0.5 M KCl (Weimberg, 1970). Any phosphate assimilated remains as orthophosphate. However, the potassium effect is reversible, and Weimberg suggests that uptake of orthophosphate and subsequent release of polyphosphate are related to changes in the conformation of the cell membrane, the probable site of action of the high concentration of K^+ . Baker's yeast, on the other hand, seems to have a specific requirement for potassium ions in the process of phosphate uptake (Schmidt et al, 1949). In fact, 0.01M potassium ions have an enhancing effect on the assimilation of orthophosphate.

Microscopy of Polyphosphate Bodies

Many organisms possess cellular inclusions which stain metachromatically with certain basic dyes (Rosenberg, 1966). These inclusions have collectively been referred to as metachromatic, volutin, or Babes-Ernst granules (Harold, 1966). The term "volutin" was first used by Meyer (1904) who noticed an accumulation of distinctive granules in Spirillum volutans. Wiame (1947 a,b; 1949) and Schmidt et al (1946) were the first to identify volutin granules as deposits of inorganic polyphosphates.

Polyphosphate bodies or granules have long been confused with other cytoplasmic inclusions. Two controversies developed based on microscopic studies and histochemical staining. Early electron microscopic studies showed that many bacteria contained granules which were highly electron scattering, and had smooth, sharply defined margins, as in Mycobacteria (Lembke and Ruska, 1940; Knaysi et al, 1951; Mudd et al, 1956), Corynebacterium diphtheriae (Morton and Anderson, 1941; Konig and Winkler, 1948; Bringmann, 1950), Staphylococcus flavocyaneus and Neisseria meningitidis (Knaysi and Mudd, 1943). Konig and Winkler (1948) first correlated the identity of these electron-scattering granules with the metachromatically staining volutin granules. This was accomplished by examining the same stained films by light microscopy and electron microscopy. Before Konig and Winkler's correlation was made, the electron-scattering granules were frequently believed to be prokaryotic nuclear bodies. This confusion arose out of light microscopy and histochemical staining, and electron microscopy.

Knaysi and Mudd (1943) and Bringmann (1950) found evidence that the electron-scattering granules contained deoxyribonucleic acid and believed they corresponded to nuclear bodies. Similar cytological evidence was presented by Lindegren (1948) who demonstrated that volutin appeared in previously phosphate-starved yeast cells on the "chromosomes" within three minutes.

From a histochemical point of view, the confusion between volutin granules and "nuclear bodies" is quite understandable. The linear array of negatively charged phosphate groups in polyphosphate bodies, and the phosphate backbone of nucleic acids are similar enough to give confusing results in histochemical staining (Fuhs, 1969). Both polyphosphates and nucleic acids are basophilic and stain metachromatically with basic dyes. Polyphosphates are more strongly basophilic, however, and retain a basic dye at a pH as low as 1.0 whereas nucleic acids destain at pH 3.5 (Fuhs, 1969). Ebel et al (1958 a) have described a technique for staining polyphosphates based on their ability to form an insoluble lead salt either at pH 3.5, or pH 1.0, depending upon the chain length of the linear polymer. Polyphosphates of chain length eight or greater are stained at either pH 3.5 or 1.0, while those with a chain length of fewer than eight retain a stain only at pH 3.5.

It had also been suggested that the electron-scattering granules in bacteria were mitochondrial equivalents. This was based on evidence by Mudd (1953, 1956), and Mudd et al (1951 a,b) that the electron-scattering granules stained intravitaly with tetrazolium salts and Janus green B. They suggested that these

granules were organized centers of oxidative-reductive activity like mitochondria and, under certain circumstances, are capable of accumulating volutin. Smith et al (1954) suggested that volutin may accumulate in more than one kind of cellular structure or inclusion.

In addition to the references already cited, polyphosphate accumulations as distinct granules have been reported by Rosenberg (1966) in Tetrahymena pyriformis by Widra and Wilburn (1959) in Aerobacter aerogenes, by Talpasayi (1963) in several blue-green algae, by Jensen (1968, 1969) in Nostoc pruniforme and Plectonema boryanum, by Stewart and Alexander (1971) in several blue-green algae, by Ebel et al (1958 b) and Keck and Stitch (1957) in a variety of organisms, and by Voelz et al (1966) in Myxococcus xanthus.

Association of polyphosphates with other cellular inclusions has been reported by several authors. Weimberg and Orton (1965) concluded that the ortho- and polyphosphates of Saccharomyces mellis are located in the protoplast and occupy a position which is different from that of phosphomono-esterase. Indge (1968) reported that in another yeast, Saccharomyces carlsbergensis, polyphosphate was located in the cell vacuole. Polyphosphates deposition in Myxococcus xanthus was found to vary in relation to the growth of the organism by Voelz et al (1966). They found three areas of deposition in the cells: 1. Dense granules around polysaccharide inclusions in the cytoplasm when phosphate in a final concentration of $5 \times 10^{-2} M$ was added to cells grown to log

phase in 5×10^{-4} M phosphate. 2. Depositions in the cytoplasm which began as dense strands, subsequently forming tightly wound bodies partially or totally surrounded by nuclear fibers, in cells which were grown in phosphate-free medium and replenished with 5×10^{-3} or 5×10^{-2} M phosphate, and 3. Dense strands scattered throughout the cytoplasm in cells grown for 20 hours without buffer, and replenished with 5×10^{-2} M phosphate.

In describing a developmental sequence of polyphosphate bodies in P. boryanum, formed under conditions of excess phosphate and continuous light, Jensen (1969) observed that all cytoplasmic inclusions are excluded from the area of polyphosphate body formation. This evidence supports that of Fuhs (1958) who demonstrated cytochemically that the polyphosphate bodies of another blue-green alga, Oscillatoria amoena, contained only polyphosphate.

The most frequently observed images of polyphosphate bodies are those of round granules, quite electron dense, and not limited by a membrane. The most thorough descriptions of polyphosphate bodies and their developmental sequences have been made by Jensen (1968, 1969). These descriptions include such features as sublimation under high electron beam intensity, also reported by Drews and Niklowitz (1957), Stewart and Alexander (1971), König and Winkler (1948), and Drews (1960). Polyphosphate bodies, due to their dense nature, often fall out or chip out in sectioning, may compress during sectioning, shrink under the electron beam, or "smear" in the direction of sectioning (Jensen 1968). The general size range of polyphosphate bodies has been reported to be between 0.1 to

2.0 μm (Stewart and Alexander, 1971; Sicko, 1972). Formation of polyphosphate bodies in Plectonema boryanum, under conditions of excess phosphate and continuous light, appears to be in the following sequence (Jensen, 1969):

1. Development of electron-lucent areas in the cytoplasm or at the cross walls,

2. Increase in size of this area to approximately the size of a mature, dense polyphosphate body,

3. Development of a porous area of medium electron density in the electron-lucent area and simultaneous deposition of polyphosphate in the cytoplasm,

4. Penetration of polyphosphate into the porous structure, resulting in the usual image of a dense granule.

This work has subsequently been confirmed by Stewart and Alexander (1971).

In addition to the descriptions previously cited, several reports have occurred in which polyphosphate is said to be deposited in morphologically different structures. Munk and Rosenberg (1969) reported that Tetrahymena pyriformis deposited polyphosphates in spherical granules which appeared to be surrounded by membranes. "Electron-scattering alveolar bodies" were reported by Fisher (1971) in a lichen phycobiont, Trebouxia erici. The appearance of these vacuolar granules varied with fixation. The polyphosphate-containing particles of Micrococcus lysodeikticus differed markedly in shape and organization from the volutin granules described in other microorganisms. Friedberg and Avigad (1968) described the electron-

dense granules ranging from 40-80 nm in diameter as part of a more complex structure. The polyphosphate bodies appeared to be organized around a granulated center in a rosettelike pattern.

Thus it can be seen from the previous discussion that the morphology and location within the cell of polyphosphate bodies seem to vary in different organisms, and under different growth conditions.

Aspects of Phosphate Metabolism

There are two distinct classes of polyphosphates found in cells: "acid-soluble", polymers which are readily extracted in cold trichloroacetic acid (TCA), and "acid-insoluble", polymers which are not extracted by cold TCA, but may be extracted by a short exposure to hot TCA (Wiame, 1949; Krishnan et al, 1957). Yoshida (1955) first reported the preparation of soluble and insoluble polyphosphates from yeast, and found that polymerization grades of acid-soluble and acid-insoluble were 10 and 50, respectively. Thus, the references in the literature are to acid-soluble forms as being short chain polyphosphates, and acid-insoluble forms as being long chain polymers (Harold, 1966; Terry and Hooper, 1970). Kanai and his associates (1963, 1965) and Miyachi and Tamiya (1961) were able further to subdivide these classes of polyphosphates into four fractions, obtained by a modified Schmidt-Thannhauser method. These fractions, obtained by successive extractions, were classified as follows:

- (1) poly-Pi "A" - cold 8% TCA

- (2) poly-Pi "B" - cold KOH at pH 9.0
- (3) poly-Pi "C" - 2NKOH, Reprecipitable by neutralizing
the extract
- (4) poly-Pi "D" - 2NKOH, soluble after neutralizing the
extract

The physiological and metabolic functions of the acid-soluble and acid-insoluble polyphosphates also seem to be distinct, and vary as a function of growth. Wiame (1949) first reported that the acid-insoluble form is metabolically more active in the cell, and that it is rapidly and reversibly transformed to orthophosphate. Katchman and Fetty (1955) substantiated this evidence in Saccharomyces cerevisiae by demonstrating that the soluble inorganic polyphosphate fraction maintained a steady-state concentration from generation to generation, while the insoluble fraction appeared only in the later stages of logarithmic growth in a nonsteady-state concentration independent of the concentration of the soluble polyphosphates. Katchman and Van Wazer (1954) postulated that the protein-complexing ability of polyphosphates differing in chain length might determine the different metabolic activities of the polyphosphates in yeast.

The differences in the amounts of the two classes of polyphosphates appear to be a function of growth. Early studies by Smith et al (1954) and Wilkinson and Duguid (1960) demonstrated that Aerobacter aerogenes contained little or no polyphosphate during exponential growth. A similar system appears to operate in Corynebacterium xerosis. Hughes and Muhammed (1962) observed an accumulation of polyphosphate in these bacteria during lag phase

after transfer to fresh medium, a decrease during the exponential phase, and an accumulation in the stationary phase. Similar results were also reported by Drews (1960), Mudd et al (1958), and Winder and Denny (1957). As a contrast to the above systems, Terry and Hooper (1970) found that short chain polyphosphates were present in constant amounts in Nitrosomonas europaea throughout growth and had a negligible turnover rate. Acid-insoluble long chain polyphosphates decreased upon transfer to fresh medium, then increased as growth proceeded and remained fairly constant. Terry and Hooper (1970) suggested that rapid hydrolysis of polyphosphate after transfer to a fresh medium was triggered primarily by the higher pH of the fresh growth medium.

It seems evident that the accumulation or degradation of polyphosphate is a function of cell metabolism. Conditions of polyphosphate accumulation under conditions of nutrient imbalance fall into two distinct patterns - (a) the "polyphosphate overplus" phenomenon as previously described, and (b) cessation of nucleic acid synthesis due to exhaustion of an essential metabolite or nutrient.

Lindegren (1948) first suggested that volutin, which Wiame (1949) had later identified as metaphosphate, was essential for cell division. Sall et al (1956, 1958) reported that in Corynebacterium diphtheriae, accumulation and disappearance of polyphosphate were physiological events related to cell division, with minimal amounts of polyphosphate present after recurrent periods of cell division. Mudd et al (1958) demonstrated that there was a competitive relationship between nucleic acid synthesis and the accumulation of poly-

phosphate in mycobacterial cells. This relationship was studied using ^{32}P and tracing the exchange from labelled polyphosphate to RNA-P. Earlier work by Schmidt et al (1956) had demonstrated that in Baker's yeast, acid-insoluble intracellular polyphosphates were utilized as efficiently as orthophosphate present in nutrient solutions for nucleic acid synthesis. Large accumulations of inorganic polyphosphates did occur in these yeast cells under conditions of complete RNA inhibition. Baker and Schmidt (1964) found that in synchronized cells of Chlorella pyrenoidosa, there was a recurrent decrease in the polyphosphate level immediately prior to and during nuclear division. This study also demonstrated that accumulations of total nucleic acid phosphorus and acid-insoluble polyphosphate phosphorus exhibited an inverse relationship to each other. Sauer et al (1969) obtained similar results with the slime mold, Physarum polycephalum. Inhibition of RNA synthesis in the plasmodium by actinomycin D resulted in a marked stimulation of ^{32}P incorporation into polyphosphate. No such correlation was found after inhibition of either DNA synthesis by 5-fluorodeoxyuridine or of protein synthesis by cycloheximide.

Evidence which does not support the hypothesis that polyphosphate-phosphorus may serve as a source of RNA-phosphorus comes from the work of Miyachi and Tamiya (1961). They demonstrated that in Chlorella ellipsoidea, the phosphorus used in the synthesis of DNA and protein was taken primarily from polyphosphates, while that used in the synthesis of RNA, phospholipid, and other polyphosphates was generally taken from an extracellular phosphorus source. This

work is supported by evidence from Kanai et al (1965) who found that poly-Pi "A" and "C" function as intermediates transferring phosphate from orthophosphate to DNA and phosphoprotein.

Extensive studies on phosphate metabolism have been made by Harold and his coworkers both on Aerobacter aerogenes and Neurospora crassa. As has been discussed before, exhaustion of an essential metabolite results in a cessation of nucleic acid synthesis. This same situation was found in A. aerogenes. When growth and nucleic acid synthesis were blocked by depriving the organisms of sulfate, assimilation of inorganic phosphate from the growth medium resulted in a slow accumulation of polyphosphate (Harold and Sylvan, 1963). This accumulation could be reversed if growth were allowed to resume. That is, polyphosphate was rapidly degraded, and the phosphate was transferred to the nucleic acid fraction. Again, evidence was accumulated for an inverse relationship between nucleic acid synthesis and polyphosphate accumulation (Harold, 1963, 1965; Harold and Harold, 1965). Using a series of mutants of A. aerogenes deficient in some aspect of phosphate metabolism, Harold (1966) has arrived at the following scheme for regulation of polyphosphate synthesis (see Table 1):

1. There is little or no deposition of polyphosphate in normally growing cells. Synthesis of nucleic acids inhibits polyphosphate synthesis and stimulates polyphosphate degradation.
2. If growth and nucleic acid synthesis are inhibited by exhaustion of an essential nutrient, polyphosphate degradation is inhibited. Possibly, the competition for ATP is

Table 1. Mutants of *Aerobacter aerogenes* deficient in polyphosphate metabolism*

Mutant	Physiological Characteristics	Enzymes Present			Enzyme Control Mechanism
		alkaline phosphatase	kinase	poly-phosphatase	
Wild	Polyphosphate not detected in growing cultures accumulates by nutrient imbalance and overplus	+	+	+	All enzymes derepressed by phosphate starvation
Pn-1	Polyphosphate accumulation by nutrient imbalance; no overplus	+	+	+	Enzymes not derepressed by phosphate starvation
Pn-2	No polyphosphate accumulation	+	-	+	Enzymes other than kinase derepressed by starvation
Pn-3	Transient accumulation of polyphosphate in growing cells	+	+	+	All constitutively elevated
Pn-4	No polyphosphate degradation	+	+	-	Enzymes other than polyphosphatase derepressed by starvation

*after F.M. Harold, 1966

relieved, and the levels of polyphosphate kinase determine the amount of polyphosphate accumulated.

3. The basis of the overplus phenomenon is elevated levels of kinase. Cells subjected to phosphate starvation are derepressed in the synthesis of kinase. Thus, exposure to inorganic phosphate results in rapid polyphosphate synthesis.

The second organism investigated by Harold was Neurospora crassa. In 1949, Houlahan and Mitchell observed an accumulation of polyphosphate in various mutants of Neurospora crassa, (Harold, 1966). Harold (1960) found that Neurospora contained high levels of polyphosphate, even during logarithmic growth. Exhaustion of a nutrient in the growth medium resulted in an accumulation of polyphosphate at the expense of RNA. During subsequent starvation, polyphosphate was degraded for RNA synthesis. Harold (1962) also found that ATP was a precursor for the overplus phenomenon, but not an intermediate in the degradation of polyphosphate.

The relationship between nucleic acids and polyphosphates is further complicated by studies which indicate that ribonucleic acid and polyphosphate may exist as a complex in several organisms. This complex was first postulated by MacFarlane (1936). RNA-polyphosphate has been demonstrated by Kulaev and Belozuskii (1958) in Aspergillus niger, by Chayen et al (1955) in Torulopsis utilis, by Winder and Denny (1957) in Mycobacteria, by Ebel et al, (1958 b, 1962) in yeast, by Correll and Tolbert (1962, 1964) and by Correll (1965)

in Anabaena and Chlorella, and by Wang and Mancini (1966) in Russell wheat. Correll and Tolbert (1962, 1964) found that in Anabaena, the complex accounted for 25-35% of the total phosphorus, and a major portion of the alga's RNA. However, there was an additional 40-50% of the total phosphorus present as uncomplexed polyphosphate. The polyphosphate-RNA complex in Chlorella was more variable; the relative amounts of the complex varied with respect to the synchronized growth cycle. Wang and Mancini (1966) isolated a RNA-polyphosphate fraction from wheat leaves. Leaves fed with ^{32}P phosphate were found to contain this complex with nearly all of the label in the polyphosphate of the complex.



It can be concluded from the previous discussion that there appears to be a correlation between levels of polyphosphate in cells and the amount of nucleic acids synthesized. In fact, there might be a direct transfer of phosphate between them, coupled with an energy prerequisite. Evidence such as this has led to the following theories of polyphosphate function.

Phosphagen Hypothesis

Much attention has been paid to the role of polyphosphate as a phosphagen. Ennor and Morrison (1958) have defined phosphagens as "those naturally-occurring phosphorylated compounds which function as stores of phosphate-bond energy from which phosphoryl groups may be transferred to ADP to form ATP as a result of enzymatic catalysis." The essential point is the direct formation of ATP at the expense of the phosphagen. Evidence exists in the literature both for and

against this hypothesis.

Kornberg (1957) and Kornberg et al (1956) described in detail an enzyme purified from Escherichia coli which reversibly catalyzed the formation of highly polymerized inorganic polyphosphate from the terminal phosphate of ATP, according to the following equations.



Polyphosphate synthesized chemically or enzymically is utilized quantitatively in the phosphorylation of ADP to ATP. Kornberg (1957) discussed the significance of this reaction and concluded that when cellular levels of ADP are high due to metabolic activity, the ADP could be phosphorylated by polyphosphate. Yoshida (1955) measured the heat of enzymic hydrolysis of insoluble polyphosphate and found it to be 10 Kcal per P-O-P linkage. Winder and Denny (1955), using cell-free extracts of Mycobacterium smegmatis, found that this extract was capable of utilizing metaphosphate for the phosphorylation of glycerol in the presence of ATP. Since their experiments were conducted under anaerobic conditions, they concluded that the metaphosphate must have provided the energy for this phosphorylation. The phototrophic Chlorobium thiosulphatophilum appears to synthesize polyphosphate when ATP is formed in excess of other requirements by photophosphorylation (Cole and Hughes, 1965). The authors also postulate that under conditions where ATP utilization is high, net breakdown of polyphosphate could occur. Purification and a study of the properties of the enzyme polyphosphate

kinase from Mycobacterium smegmatis (Suzuki et al, 1972) support these earlier results.

Evidence contradicting the role of polyphosphate as a phosphagen is discussed in detail by Harold (1966). Two lines of research are responsible for this interpretation: (1) Harold (1962) has demonstrated that in intact cells of Neurospora crassa, polyphosphate does not break down if energy generation is limited or blocked. Similar evidence is provided by Kaltwasser (1962) for Hydrogenomonas. Polyphosphate breakdown can occur in yeast treated with iodoacetate, but the pathway is not through conversion to ATP, (Langen and Liss, 1958; Langen, 1965). (2) In Aerobacter aerogenes, there is direct evidence that polyphosphate degradation is hydrolytic, resulting in dissipation of the energy-rich bond (Harold and Harold, 1965). From this evidence, Harold (1966) concludes that the hypothesis of polyphosphate as a phosphagen is not tenable.

Polyphosphate as a Phosphorus Reserve

Much evidence has been presented to support the theory that polyphosphate is a storage form of phosphorus. As has already been discussed, insoluble polyphosphate can serve as a source of phosphorus in nucleic acid synthesis, most often RNA synthesis. Polyphosphate, due to its structure, minimizes disturbance of osmotic equilibrium in the cell, and thus makes it an attractive compound for storage in the cell (Voelz et al, 1966; Harold, 1966). As will be discussed in a later section, the concentration of phosphorus in the environment is usually low. If the scheme proposed by Harold as discussed previously were operative, cells subjected to low concentrations of

phosphorus would have elevated levels of polyphosphate kinase, the enzyme responsible for polymerization of orthophosphate. Thus, when the organisms come into contact with a phosphate source, the organism would be able to accumulate a phosphorus reserve for later synthesis of cellular compounds (Harold, 1966). This theory is also substantiated by Kaltwasser (1962) who found similar results in Hydrogenomonas.

Ecological Aspects of Phosphorus in Natural Waters

The role of phosphorus in the eutrophication process has long been a controversial subject. Nutrients such as carbon, nitrogen, and phosphorus are often referred to as "limiting" or regulating nutrients in an aquatic eco-system, limiting in the sense that they control photosynthetic productivity. In fact, nutrient concentrations of these and other elements are often used to classify lakes into either oligotrophic or eutrophic categories (Rodhe, 1969).

Eutrophication, in the most popular definitions, is usually man-centered, and refers to an enrichment process which is vastly accelerated by man (Hasler, 1947; Likens, 1972). From an ecosystem point of view, eutrophication may be broadly defined as follows (Likens, 1972):

"Eutrophication...nutrient or organic matter enrichment, or both, that results in high biological productivity, and a decreased volume within an ecosystem."

Likens (1972) also points out that eutrophication is often considered as a form of pollution, but the two terms are actually not synonymous.

Phosphorus, especially in the form supplied by effluents con-

taining synthetic detergents, is usually implicated as the main source of nutrient enrichment responsible for accelerating eutrophication. This topic, as will be reviewed, is quite extensive, and yet the role of phosphorus, as it involves organisms in an aquatic ecosystem, is not clearly defined.

Nutrient Sources in Aquatic Ecosystems

Phosphorus can be found in a variety of forms in water. The sources of phosphorus-containing compounds are equally diverse (Table 2). It has been estimated that millions of pounds of phosphates per year are contributed from these sources (Ferguson, 1968). The phosphate source most frequently cited as the principle factor causing blooms of algae is synthetic detergent builders, consisting for the most part, of sodium tripolyphosphate.

Evidence exists again, for both sides of this question. An important factor lies in the definition of a "limiting" nutrient, and changes in the concentration of this nutrient. Gibson (1971) reviews the concept of nutrient limitation from the following three definitions:

1. An organism is limited when it is not growing as fast as it is theoretically able to.
2. A factor is said to be limiting when it is in such short supply that no growth is possible.
3. A factor is not limiting, if when it is increased, no effect on growth is observed.

The third definition is most commonly encountered, and evidence both for and against phosphate being a limiting nutrient is approached

TABLE 2

SOURCES OF NUTRIENTS OCCURRING IN NATURAL WATER (Modified from P. J. Weaver, 1969; L. I. Keup, 1968)

	<u>SOURCE</u>	<u>TYPE OF PHOSPHORUS</u>	<u>AMOUNT</u>	<u>REFERENCES</u>
I	Municipal Treatment Plants and Private Waste Disposal Systems.	total soluble P output for all types of sewage	1-13mg/l	Fitzgerald and Rohlich (1958) Lewin (1973)
	A. Domestic		10mg/l	Mitchell (1971)
	1. Human Wastes	Ortho P-P		
	2. Soaps and Detergents	Organic	total estimated all sources	
	3. Household food	(2) condensed inorg. PO ₄	3 lb. mean per capita per annum	Bush and Mulford (1954) Devey and Harkness (1973)
	B. Industrial			
II	Industrial Wastes Discharged directly to Waterways			Solt (1973)
III	Water Treatment Chemicals			
IV	Land Runoff			
	A. Urban Runoff and Drainage			
	B. Stormwater	measured as P	0.82 lbs. PO ₄ -P/yr/acre, 10-1400ug/l	Weibel <u>et al.</u> , (1966) Sylvester (1961)

<u>SOURCE</u>	<u>TYPE OF PHOSPHORUS</u>	<u>AMOUNT</u>	<u>REFERENCES</u>
C. Rural Runoff and Drainage			
1. Agricultural			
a. soils (erosion)	measured as P	13 lbs./acre	Fippen (1945)
b. fertilizers	slurries containing P		Cooke and Williams (1973)
c. animal excrement		up to 1.5mg/l	Cooke and Williams (1973)
2. Non-agricultural land			
a. decaying leaves	orthophosphate	54-230ugP/gram leaves	Cowan and Lee(1971)
b. wild animal wastes			
V Groundwater	PO ₄ -P	10-70ug/l	
VI Reserves in Lakes			
A. Bottom Muds	Organic, PO ₄ -P		Golterman (1973)
B. Living Aquatic Organisms			
C. Lake Water	PO ₄ -P	up to 1 mg/l in peat-rich water	Golterman (1973)
D. Peat			
VII Atmosphere			
A. Rainfall	not determined	0-49ug/l 0-80ug/l	Hutchinson (1957) Wiebel <u>et al</u> (1966)
B. Dustfall			

from this viewpoint.

Phosphorus has long been found to be an important minimum factor for plant growth in natural waters (Sawyer, 1947; 1952; Ohle, 1953; Fuhs et al, 1972). Thomas (1953) observed that phosphorus and nitrogen were minimum factors for algal growth. The high phosphorus content of waste waters was a key factor in the acceleration of eutrophication above its natural rate. If the phosphorus demand of the algae had been met, nitrogen replaced phosphorus as the minimum factor. Fuhs et al (1972) also observed multiple nutrient limitation in Lake George, New York. They found that nitrogen and phosphorus could act as limiting nutrients, either simultaneously or alternating, with time and space. With nitrogen limitation in effect, low concentrations of phosphorus could not be interpreted as indicating phosphorus limitation, because nitrogen limitation favored luxury uptake of phosphorus. Schelskè and Stoermer (1972) found that in natural phytoplankton assemblages enclosed in plastic bags in Lake Michigan, phosphorus was the limiting nutrient. This resulted in lower concentrations of silica in the lake, and replacement of diatoms with nonsiliceous forms, such as blue-green and green algae. Edmunson (1961, 1969, 1970, 1972) demonstrated that the phytoplankton population of Lake Washington (Seattle) was dependent upon the increasing volumes of effluent from secondary sewage treatment plants. There was a strong correlation between the abundance of phytoplankton in the summer, and high levels of phosphate, but not nitrate or carbon dioxide, in the water during the winter months. Diversion of the effluents resulted in a

decreased phosphate concentration, and a reversal of cultural eutrophication in Lake Washington. In fact, by the time half the effluent had been diverted, the phosphate concentration started to decrease. Edmunson (1972) then made predictions about the possible changes that could occur. Since about one-half of the phosphate in the sewage came from detergents, elimination of all detergents from sewage should lead to the same results as following diversion of one-half of the effluent. The lake could then tolerate a much larger human population without their accompanying detergents.

Similar results in determining in situ effects of added phosphates in natural waters were obtained by Powers et al (1972). Lakes of varying productivity in both Minnesota and Oregon were studied, and it was found that phosphorus appeared to be the primary controlling nutrient in enrichment experiments. They postulated that several lakes could be restored by the removal of phosphorus from municipal waste by advanced treatment methods. Sonzogni and Lee (1972) also demonstrated that effluent diverted from a number of Wisconsin lakes resulted in a reduction of phosphorus content of the lakes, as well as a decrease in the frequency and severity of blue-green algal blooms. Pitcairn and Hawkes (1973) demonstrated that there was a general positive correlation between the standing crop of Cladophora and the phosphorus concentration of several river waters. They confirmed the importance of phosphorus by showing that the growth of Cladophora in waters upstream of sewage discharges could be increased to downstream levels by the addition of phosphorus.

Taylor (1967) reviewed the phosphorus concentrations that limit algal growth. Sawyer (1952) found that phosphorus concentrations below 0.01 parts per million severely limit algal growth, while concentrations of 0.05 parts per million or higher permit profuse growth. Most uncontaminated lakes contain between 0.01 and 0.03 parts per million of phosphorus; thus, adding a relatively small amount of phosphorus to the lakes, to increase the level to 0.05 ppm, is likely dramatically to increase productivity (Sawyer, 1952).

This view is not supported by a number of workers. Kuentzel (1969) reviewed the carbon dioxide-phosphate controversy and concluded that CO_2 and organic matter supporting growth of bacteria were responsible for massive algal blooms rather than phosphorus. He suggested that a symbiotic relationship between bacteria and algae could open up another possibility for reductions in algal growth via control of bacteria. The line of succession suggested was: organic matter \longrightarrow bacteria \longrightarrow CO_2 \longrightarrow algae. This review was attacked heavily by Shapiro (1970), who pointed out the fact that CO_2 is not limiting, and in spite of high concentrations of CO_2 , algae are not able to grow unless there is a sufficient supply of phosphorus.

Mitchell (1971) found that the eutrophication potentials of a phosphate-containing detergent and two phosphate-free detergents were not significantly different. He pointed out that elimination of phosphates in domestic sewage due to detergents would still result in a concentration of 3-4mg P/l, using conventional primary

water treatment. He did not consider the use of activated sludges. His estimates of phosphate concentrations in domestic sewage are in agreement with those of Hudson and Marson (1970), and Marson (1971). However, Marson reported up to 90% phosphorus removal by any of the precipitation processes currently available, and 80% by luxury consumption by activated sludge.

Levin and Shapiro (1965) demonstrated that luxury uptake of orthophosphate by sludge organisms occurred in the absence of growth. Uptake was dependent upon the dissolved oxygen and pH of the mixed liquor, with maximum uptake occurring in the pH range 7.0-8.0. Shapiro (1967) later determined that settling of activated sludge in settling ponds of water treatment plants led to anoxia of the organisms, and subsequent release of the phosphate. Yall et al (1970) supported the concept of biological luxury uptake by activated sludges using both radioactive tracers and inhibitor studies.

Jenkins et al (1973) studied the environmental impact of detergent builders in California waters and found that detergents accounted for about 35% of the total phosphorus released to surface waters. They suggested that the algae in the California waters did not appear to be phosphorus-limited, so that control of phosphorus input would do little to control eutrophication. If phosphate control were necessary, they suggested point-source elimination for a particular area and not solely detergent phosphate elimination.

The controversy of phosphate removal still exists. The problem

is further complicated by the fact that synthetic detergent builders are condensed phosphates, usually sodium tripolyphosphate (Davis and Wilcomb, 1967, 1968). The studies discussed so far were concerned with utilization of orthophosphate in natural waters by phytoplankton. Thus, two more problems are involved: 1. The natural hydrolysis of condensed phosphates in natural waters, and 2. uptake and metabolism of condensed phosphates by phytoplankton.

Hydrolysis of Condensed Phosphates

Synthetic detergent builders, condensed phosphates, appear in sewage effluents and receiving waters. These compounds, under various conditions, have a tendency to react with water and ultimately form orthophosphate. Engelbrecht and Morgan (1959) found that sodium tripolyphosphate and tetrasodium polyphosphate were subject to degradation in natural waters, the hydrolysis rates varying with the water sample. They also indicated that biological life exerted an effect on the rate of degradation, since filtration of raw water decreased the rate of degradation.

Clesceri and Lee (1965a,b) studied the rates of hydrolysis of condensed phosphates in non-sterile and sterile environments. They found that pyrophosphate and tripolyphosphate were apparently not as available a phosphorus source as orthophosphate. Using both unialgal and axenic cultures of Chlorella, the condensed phosphate compounds were completely hydrolyzed in a short time by the unialgal cultures, but not by the axenic cultures. They attributed the increased hydrolysis rates to the presence of non-algal microorganisms in the unialgal culture. Clesceri and Lee (1965b)

also showed that the rate of hydrolysis of condensed phosphates was higher by several orders of magnitude in sterile lake water and algal culture media than in distilled water at a similar pH and temperature. Hydrolysis rates of both pyrophosphate and tri-polyphosphate were highest in those solutions containing the highest concentrations of calcium ion. When comparing these rates to those determined in the presence of microorganisms under similar conditions (Clesceri and Lee, 1965a) it was found that enzymatic processes controlled the aqueous environmental chemistry of the condensed phosphates.

Clesceri and Lee (1965a) summarized the studies on the factors that influence the rate of condensed phosphate hydrolysis as follows:

<u>Factor</u>	<u>Effect on rate</u>
A. Temperature	10^5 - 10^6 x faster from freezing to boiling.
B. pH	Rates of hydrolysis of all condensed phosphates are higher in acidic media; tripolyphosphate hydrolysis can be base-catalyzed, but it is most stable in the pH 9-10 range.
C. Enzymes	Divalent cations in combination with enzymes accelerate hydrolysis, especially magnesium; adaptive enzymes, usually phosphatases, are produced by many organisms.

- D. Colloidal gels Hydrated oxides of iron, cobalt, nickel aluminum, and rare earths accelerate hydrolysis.
- E. Complexing cations pH effect is absent when there are no cations; calcium increases hydrolysis rates more than sodium.
- F. Concentration Hydrolysis of condensed phosphates are first order processes; the rates are proportional to the concentration.
- G. Independence of hydrolysis in a mixture of phosphates Hydrolysis of different species in the same dilute solution proceeds independently.

The data presented by Clesceri and Lee (1965a,b) were later substantiated by Davis and Wilcomb (1967, 1968). Cultures of several green algae, grown in nutrient medium with phosphate concentrations close to those expected in sewage, were capable of degrading polyphosphates to orthophosphates. Environmental parameters affected orthophosphate assimilation. Axenic cultures of the algae demonstrated hydrolytic ability, a greater hydrolytic capacity existing in moving systems than in static ones. Davis and Wilcomb (1968) also demonstrated that several genera and species of blue-green algae were able to utilize condensed phosphates, and, in fact, return them to the aqueous environment during certain growth phases. Again, uptake of the condensed phosphates by the blue-greens was greater in moving columns than in static cultures.

It has already been suggested (Clesceri and Lee, 1965a) that

microbial activity is responsible for the degradation of condensed phosphates to a much greater extent than other environmental parameters. This microbial activity appears to be the result of phosphatases, enzymes which catalyze the release of bound phosphates. Hydrolysis of tripolyphosphate results in a unit of pyrophosphate and a unit of orthophosphate, whereas hydrolysis of pyrophosphate yields two units of orthophosphate (Clesceri and Lee, 1965a). Kornberg (1956) isolated tripolyphosphatase, pyrophosphatase, and trimetaphosphatase from yeast cells. Eppley (1962) found that orthophosphate was released when living pieces of Porphyra were incubated with several condensed phosphates and ATP. Eppley (1962) indicated enzymatic catalysis of these compounds, and also concluded that both calcium and magnesium were essential for hydrolysis.

The presence of adaptive phosphatases has been observed in a variety of organisms. Galloway and Krauss (1963) found an adaptive pyrophosphatase associated with the cell wall of Chlorella. Overbeck (1961a,b) found intercellular phosphatase activity in Scenedesmus quadricauda when the substrates were condensed or organically bound phosphates. A number of correlations have been made numerous times and in a variety of organisms between the levels of alkaline phosphatase and the amount of phosphate in the growth medium. Alkaline phosphatase appears to be induced when the external phosphate source is limiting in E. coli (Torriani, 1960), B. subtilis (Cashel and Freese, 1964), Vibrio parahaemolyticus (Sakaguchi et al, 1972), Anacystis nidulans (Reichardt, 1971), and Anabaena flos-aquae (Bone, 1971), and a variety of algae (Fitzgerald

and Nelson, 1966). Torriani (1960) suggested that it is a means of obtaining phosphate from organic phosphate when the supply or concentration of orthophosphate becomes limiting in the medium.

MATERIALS AND METHODS

Growth Curves

The test organism was chosen after series of growth curves in various culture media were determined. The two blue-greens initially selected were Plectonema boryanum Gomont (Indiana Culture Collection No. 581) and Oscillatoria tenuis Ag. (Indiana Culture Collection No. 428), obtained from the Starr Culture Collection (Starr, 1964). This selection was based on their ability to grow well in a defined medium, Modified Fitzgerald (Fitzgerald et al, 1952; Zehnder and Gorham, 1960). See appendix A for composition of the medium. Growth curves for the algae were determined by inoculating a known dry weight of the algae into sterile culture tubes containing 20 ml of Modified Fitzgerald's medium. The tubes were then placed in a Sherer-Gillette growth chamber adjusted to 1. 500 ft-candles of illumination (5330 lux) from incandescent and fluorescent sources, 2. 25°C and, 3. an alternating 12 hour day/night cycle. Growth was monitored for a period of 28 days as a function of increase in dry weight per unit volume. Stock cultures were also maintained under these conditions for one month, and subsequently placed under 200 ft-candles of illumination. Transfers to fresh medium were made from cultures which were between two and four months old.

Dry Weight Determinations

Dry weight of the samples was found to be the most consistent and easiest method for determining growth of the algae. Both blue-greens tested are filamentous and have sheaths which make dispersal and rupture of filaments into individual cells difficult for counting or spectrophotometric analysis. Millipore filters, type HA with a pore diameter of 0.45 μ m were predried by heating to 100°C for 24 hours, and weighed after cooling to determine the dry weight. Known volumes of algal suspensions were then syringed sequentially ten times each through 14, 18 and 22 gauge sterile disposable needles, and then passed through the predried Millipore filters. After drying the filter under an incandescent lamp, the filters were then dried again under the previously described conditions, and weighed when cool. The difference in the weight of the filters was taken to be the dry weight of the algae per unit volume filtered. Normally, a 10 to 20 ml aliquot containing between 200 and 1000 mg algae per liter was filtered.

Correlation of Dry Weight to Cell Counts

Although dry weight analysis was chosen to be the method for quantitating the algal suspensions, correlations were made between actual cell number and dry weight. This was accomplished by extensive syringing of the cultures through 22 gauge needles and recording the following information using a hemocytometer and reticule: 1. Number of algal filaments per unit volume of culture, 2. Length of individual filaments, and 3. Number of cells per filament. In this manner, the number of cells per unit volume was

determined, and correlated to a dry weight analysis of that culture. Analyses of cell number, dry weight, and average cell size were made for logarithmic phase culture, phosphate starved cultures, and cultures at the end of the four hour rapid uptake period.

Phosphate Assay and Calibration

Phosphate depletion of the medium was also monitored during all growth determinations. This was accomplished by testing the growth medium without the algae. In all cases, the algae were removed by Millipore filtration as previously described. A known volume of the filtrate was diluted volumetrically so that the spectrophotometric absorbance values were in the range 0.3 to 0.7 whenever possible. The two tests used for the determination of orthophosphate were the stannous chloride method (American Publ. Health Assoc., 1965) and the single solution method of Murphy and Riley (1962). See Appendix B for the composition of the solutions.

The method selected to be the most reliable was that of Murphy and Riley (1962). The orthophosphate determinations are colorimetric tests; the absorbance of the phosphomolybdate complex formed is a function of orthophosphate concentration in solution. The colored complex formed during the Murphy-Riley determination is stable for a period of 24 hours, and there is less interference due to arsenic or salts. The stability of the colored complex formed in the stannous-chloride method is considerably less. Forty ml of the solution to be tested were placed in a 50 ml graduated cylinder. Eight ml of the mixed reagent, and two ml of glass distilled water were added so that the final volume of the solution was fifty ml.

The solution was then mixed thoroughly and color development was allowed to proceed from at least 10 minutes up to a period of 1 hour. Percent transmittance was read directly and converted to an absorbance value. All readings were taken at 880 nm on a Spectronic 20 spectrophotometer equipped with infrared sensitive phototubes and filters. Calibration curves were used to determine the sensitivity of the test. The absorbance value divided by the slope of the calibration curve gave a direct reading of phosphorus concentration as phosphate. All phosphate determinations were carried out in glassware which was acid washed and used only for these tests.

Calibration curves were determined in the following manner: A standard phosphate solution containing 0.1757 grams of potassium dihydrogen phosphate per liter was prepared. This solution contains 40 mg P (as phosphate) per liter. Dilutions of the stock solution were made to the following concentrations:

80ug P/l	(3.2ug P/40 ml)
160ug P/l	(6.4ug P/40 ml)
320ug P/l	(12.8ug P/40 ml)
400ug P/l	(16.0ug P/40 ml)
640ug P/l	(25.6ug P/40 ml)

The percent transmittance values were determined, and converted to absorbance. The values were measured at 880 nm in 2.56cm cells. Beer's law is obeyed in this concentration range, and the method gives an excellent reproducibility.

Starvation Conditions

Cells of Plectonema boryanum were grown in Modified Fitzgerald's

medium containing 8 - 10 mg PO_4 per liter for 14 - 17 days at 25°C, 500 ft-candles of illumination, and an alternating 12 hour day/night cycle. To induce phosphorus starvation, cultures were harvested aseptically by centrifugation in a Sorvall refrigerated centrifuge at 21°C and 12,100 xg for 10 minutes. The medium containing phosphate was decanted, and the cells were washed 3 times in sterile medium free of phosphate. After the final centrifugation, the cells were resuspended in phosphate free medium and then placed under the original environmental conditions. Optimal starvation conditions were determined by prolonging starvation up to a length of 12 days.

"Overplus" or Rapid Uptake

Luxury storage of phosphate was induced by starving the algae of phosphate for 5 days, and then inoculating into medium containing a known amount of phosphate. This phosphate concentration was varied in the range 0.1 to 112 mg PO_4 per liter. The rapid uptake was monitored in one or more of several ways: 1. increase in dry weight of the algae, 2. depletion of phosphate from the medium, 3. increase in total phosphate in the cells, 4. increase in various phosphate-containing fractions, or 5. light or electron microscopy of the samples.

Inhibitors and Temperature Studies

For uptake studies involving inhibitors or temperatures other than 25°C, the algae were treated in the following manner. At the end of the starvation period, the algae were pelleted by centrifugation and resuspended in medium containing no phosphate and containing the appropriate concentration of inhibitor. The algae were then

placed under the appropriate growth conditions and incubated with the inhibitor for a period of 1 hour prior to the addition of phosphate in the normal uptake manner. The three inhibitors tested were 2, 4-dinitrophenol ($5 \times 10^{-3} \text{M}$), mercuric chloride ($1 \times 10^{-2} \text{M}$), and sodium fluoride ($1 \times 10^{-3} \text{M}$). Uptake studies involving two different temperatures were treated somewhat differently, depending on the temperature. For uptake at 37°C , the algae were first acclimated to that temperature for a period of one month, then logarithmic phase cells were starved in the normal manner. This was possible since the algae grow well at this temperature. Uptake at low temperatures involved the growth of the algae at 25°C , and preincubation of the algae at 4°C for 1 hour prior to uptake. This procedure was followed because there is virtually no growth at this low temperature.

Total Phosphate Determinations

Batterton and Van Baalen's (1968) modification of the Menzil-Corwin (1965) potassium persulfate ($\text{K}_2\text{S}_2\text{O}_8$) digestion was used for the assay of total phosphorus levels in the cells. A known aliquot of washed cell suspension, containing no more than 1 g algae per liter was placed in screw cap culture tubes containing 0.5 grams of potassium persulfate. The tubes were then sealed loosely and autoclaved at 121°C for thirty minutes. The clear contents of the tube were transferred quantitatively and assayed for orthophosphate by the Murphy-Riley technique. Total phosphorus was reported as ug P per ug dry weight algae.

Extraction Scheme for Polyphosphates

The extraction method chosen was Harold's (1960, 1963) modification of the Schmidt-Thannhauser extraction scheme for nucleic acids (1945). Both trichloroacetic acid (TCA) and perchloric acid were tested initially. TCA was found to be more reproducible, and was chosen subsequently for all extractions. A flow chart of this scheme is presented in Figure 2.

A. Cold TCA Extraction

Washed algal suspensions were extracted twice with 5 ml of cold (4°C) 5% TCA. The supernates were pooled along with the distilled water wash of the residue. This supernate was assayed both for orthophosphate and total phosphate. In all cases, the residue was separated from the supernate by centrifugation in a VWR MSE GT 2 centrifuge with a swinging bucket rotor at 3280 rpm (1745 xg) for 10 minutes.


B. Lipid Extraction

The residue of the cold TCA extraction was then extracted with 5 ml of absolute ethanol at room temperature for 30 minutes, followed by extraction in 5 ml of ethanol-ethyl ether (3:1). The mixture was boiled for 1 minute, and left to stand at room temperature for 20 minutes. The two supernates were pooled along with a distilled water rinse of the residue and assayed both for orthophosphate and total phosphate.

C. Hot TCA Extraction

The lipid-free residue was then dried by heating the test tubes in a 70°C water bath, and then extracted twice with 5 ml portions of

FRACTIONATION SCHEME FOR THE EXTRACTION OF PHOSPHORUS-CONTAINING COMPOUNDS



5% TCA warmed to 70°C at 15 minute intervals. The supernates were pooled along with a distilled water rinse of the residue and assayed both for orthophosphate and total phosphate. The residue remaining after this extraction was then digested with potassium persulfate as previously described, and then analyzed for orthophosphate content.

D. Separation of Nucleic Acids

200 mg of sulfuric acid washed Norit A was added per 5 ml of extract. Nucleic acids were then eluted with alcoholic ammonia from the charcoal, and either analyzed for total phosphorus concentration, or the extinction at 260 nm was determined.

For all total extractions, the total phosphate content of the algal suspension was determined before equivalent aliquots were fractionated. In this manner, percent recovery could be calculated. Secondary precautions were also taken by total phosphate determinations on residues after each extraction, and by processing triplicate samples in all cases.

Electron Microscopy

Algae were harvested from cultures grown under the various culture conditions previously described, and prepared for electron microscopy in the following manner. Algae were pelleted from the original medium by centrifugation at 1745 xg. The supernatant medium was discarded and the algae resuspended in Modified Kellenberger fixative according to the method of Pankratz and Bowen (1963). The fixative employed was 1% OsO₄ in Michaelis buffer at pH 6.2 for 3 hours at room temperature. The algae were then dehydrated in a

graded ethanol series and embedded in Epon 812 according to the method of Luft (1961). See Appendix C for fixation and embedding details.

Sections approximately 500 A thick were cut with a DuPont diamond knife on a LKB Ultratome III, and collected on clean 300 mesh copper grids. The sections were post stained with saturated uranyl acetate in methanol (Stempak and Ward, 1964) for 10 minutes, or with lead salts (Reynolds, 1963), separately or in combination with uranyl salts, and examined in an Hitachi HU 11E electron microscope operating at 75 kV. Pictures were taken on Kodak Contrast Projection Plates and developed with D-19 diluted with 1 part water at 68°F for two minutes.

Light Microscopy

Algal suspensions were stained for polyphosphates by the method of Ebel et al (1958) and Jensen (1968). This staining procedure employs the ability of lead salts to complex with polyphosphate and remain stable at low pH values. The algae were initially fixed in 3% glutaraldehyde in 0.1 M cacodylate buffer adjusted to pH 6.2 for a period of 1 hour at 4°C. The algae were then rinsed five times in 0.1 M cacodylate buffer, pH 6.2, and incubated in 20% lead nitrate, adjusted to pH 3.4 with acetic acid, for 4 hours at room temperature. After the incubation in lead nitrate, the algae were washed thoroughly five times with distilled water, then placed in 1% ammonium sulfide at room temperature for ½ hour. They were again washed with distilled water and then examined with a Zeiss light microscope (Jensen, 1968). Pictures

were taken either with Polaroid P/N 55 film, or Kodak Tri-X film and developed in Microdol X diluted with 1 part water.

X-Ray Energy Dispersive Analysis

Sections, approximately 0.5 μ m in thickness, were cut on glass knives and mounted on 200 mesh copper grids. The sections were then examined and analyzed in a JEM-100B analytical electron microscope fitted with a scanning attachment, a $\pm 60^\circ$ side entry goniometer stage, and either EDAX or KEVEX energy dispersive x-ray Si (Li) detector having 180 eV resolution. For orientation purposes, the sections were examined at either 80 or 100 kV in the transmission or scanning modes. The areas to be analyzed were then selected, and the accelerating voltage reduced to 40 kV for elemental analysis. The beam spot could be reduced to less than 50A in diameter, much smaller than the "target" area of this study, so the beam was in all cases reduced to the size of the polyphosphate body, or a comparable area if no polyphosphate body was present.

All analyses were carried out in one of two ways: either total counts of emitted x-rays were read for a fixed time period, or total counts, an arbitrary number, and regardless of time, were read. The fixed time was usually 100 seconds, and the fixed count was usually 20,000.

Samples examined were treated in three ways:

1. Glutaraldehyde - fixation and Epon embedding,
2. Osmium - fixation and Epon embedding, and
3. Osmium - fixation and Durcupan embedding. Durcupan is a water-soluble embedding medium. See Appendix 2, Schedule B for the

procedure of Durcupan embedding. All samples examined were starved of phosphate for 5 days, and fixed after 4 hours of rapid uptake.

Determination of Standard Deviation and Standard Error

The standard deviation for the physiological experiments was determined for all experiments where at least four samples were run.

The standard deviation was calculated as follows:

1. The arithmetical mean (\bar{x}) was determined by adding all of the individual observations (X_i) and dividing the total by the number of individual samples (N).
2. The individual deviation from the mean ($X_i - \bar{X}$) was calculated.
3. Each individual deviation from the mean was squared ($(X_i - \bar{X})^2$) so that positive values were obtained.
4. The sum of the squared deviations from the mean was determined ($\sum (X_i - \bar{X})^2$).
5. The standard deviation of the sample was calculated using the formula.

$$S = \sqrt{\frac{\sum (X_i - \bar{X})^2}{N - 1}}$$

The standard deviation enables one to measure the possible error of the estimate. It is also possible to measure the sample mean by an interval. The value $\frac{s}{n}$ is the standard error of the sample mean.

In figures showing the results in graphic form, the confidence interval is represented by uncolored circles superimposed on each bar (Machlis, L., and J. G. Torrey. Plants in Action. W. H. Freeman and Company, San Francisco, 1956, pp. 111-145).

RESULTS AND OBSERVATIONS

Growth Curves

The test organism, Plectonema boryanum, was chosen after a series of growth curves were determined both for P. boryanum and Oscillatoria tenuis. P. boryanum is able to grow well with phosphate concentrations ranging from 0.8 mg PO₄ to 1100 mg PO₄ per liter (Figures 3 - 9). Growth of P. boryanum was also monitored simultaneously for phosphate depletion of the medium by the alga. In all phosphate concentrations monitored, the phosphate remaining in the medium was reduced substantially the first day after transfer, rose slightly on the third or fourth day, and then decreased gradually through the remainder of the 28 day period (refer to Figures 3 - 9).

The growth curves varied somewhat with the different concentrations. All growth curves were characterized by an increase in dry weight, often more than four-fold, on the first day after transfer. This initial peak then diminished during the lag period. The growth of Plectonema boryanum is sigmoidal; lag, log, stationary, and decline phases are recognizable. The length of the lag and log periods seem to be a function of the phosphate concentration in the medium. Longer log periods of growth are encountered at higher phosphate concentrations (Figures 7 and 8). Lag period appears to be longer at lower phosphate concentrations.


Figures 3-6. Growth curves of Plectonema boryanum and phosphate depletion of culture medium. All growth curves were determined at 25°C, 500 ft-candles of illumination, and a 12 hour alternating day/night cycle. Growth is represented by circles and phosphate depletion by squares. Each reading represents the mean of these determinations.

Figure 3 - 1 mg PO₄ / liter culture medium

Figure 4 - 2 mg PO₄ / liter culture medium

Figure 5 - 5 mg PO₄ / liter culture medium

Figure 6 - 8 mg PO₄ / liter culture medium




Figures 7-9. Growth curves of Plectonema boryanum and phosphate depletion of medium. All growth curves were determined at 500 ft-candles of illumination, 25°C, and a 12 hour alternating day/night cycle. Growth is represented by circles and phosphate depletion of the medium by squares.

Figure 7. 10 mg PO₄/liter

Figure 8. 100 mg PO₄/liter

Figure 9. 1000 mg PO₄/ liter



This data led to routine culture of P. boryanum at either 8 or 10 mg of phosphate per liter. At these concentrations, log phase occurs approximately between 13 and 17 days of culture (Figures 6 and 7). These are also normal phosphate levels for synthetic media.

Oscillatoria tenuis was also cultured at the same phosphate concentrations as P. boryanum, but with quite different results (Figures 10 - 16). O. tenuis did not grow well in phosphate concentrations above 10 mg PO_4 per liter (Figures 15 and 16). At 1000 mg PO_4 per liter, O. tenuis increased in dry weight two days after transfer. However, the culture was not able to survive at this high phosphate concentration for more than four days (Figure 16). The phosphate depletion from the medium was similar to that of P. boryanum (Figures 10 - 16). Plectonema boryanum was then selected for all subsequent experiments due to its ability to tolerate a wide range of phosphate concentrations.

Figure 17 demonstrates the growth of the alga at 37°C. This rate was determined so that the alga could be acclimated to that temperature for uptake studies. By raising the temperature 12°C above room temperature, the growth rate was approximately doubled so that log phase occurred at 7 days. The 10 mg of phosphate initially present in the medium was depleted after 1 day of culture, and remained so for the 14 day period.

Total Phosphate Present as a Function of Growth

The initial increase in dry weight of the alga upon transfer to fresh medium was of particular interest. It was initially thought to be an artifact, but it was also accompanied by a tremendous loss


Figures 10-13. Growth curves of Oscillatoria tenuis and phosphate depletion of the culture medium. All growth curves were determined at 25°C, 500 ft-candles of illumination, and a 12 hour day/night cycle. Growth is represented by circles and phosphate depletion by squares.

Figure 10. 1 mg PO₄ / liter culture medium

Figure 11. 2 mg PO₄ / liter culture medium

Figure 12. 5 mg PO₄ / liter culture medium

Figure 13. 8 mg PO₄ / liter culture medium



Figures 14-16. Growth curves of Oscillatoria tenuis and phosphate depletion of the culture medium. All growth curves were determined at 25°C, 500 ft-candles of illumination, and a 12 hour alternating day/night cycle. Growth is represented by circles and phosphate depletion of the medium by squares.

Figure 14. 10 mg PO₄ / liter culture medium

Figure 15. 100 mg PO₄ / liter culture medium

Figure 16. 1000 mg PO₄ / liter culture medium





Figure 17. Growth curve of Plectonema boryanum and phosphate depletion of the culture medium. Growth conditions were 10 mg PO_4 / liter culture medium, 37°C , 500 ft-candles of illumination, and an alternating 12 hour day/night cycle. Growth is represented by circles and phosphate depletion of the culture medium by squares.



of phosphate from the medium and was present during all growth determinations. The total cell phosphorus was then measured for cultures of the alga between 1 and 51 weeks old. These results are presented in Figure 18. P. boryanum, inoculated into fresh Modified Fitzgerald containing between 8 and 10 mg PO₄ per liter medium, reaches a peak in total cell phosphorus at approximately 1 month of culture. This value was 53 ug P per mg dry weight algae (Figure 18). The cell phosphorus then declined and began to attain a constant value at about 2 months of culture; this lower limit of phosphorus concentration being approximately 4 ug P per mg dry weight algae (Figure 18). Thus, transfers to fresh medium were always made from cultures which had reduced total cell phosphorus levels.

Similar values for total cell phosphorus levels were also determined for the 37°C cultures (Figure 19). It was found that the highest levels occurred on the seventh day after transfer of the cultures. This value was 14 ug P per mg dry weight algae. The total cell phosphorus then declines to approximately 3.5 ug P per mg dry weight algae at the end of 34 days in culture. The lower values for the total cell phosphorus are similar for both the 25 and 37° cultures, while the maximum values are considerably higher in the 25°C cultures (Figures 18 and 19).

Determination of Optimum Starvation Length for Rapid Uptake

for All the Following Experiments

Logarithmic phase cells were used for all following experiments. Cells in this metabolic state were chosen because the amount of

Figure 18. Total Phosphorus Content of Plectonema boryanum as a function of culture age. The distribution is for cultures maintained at 25°C, and under the conditions as listed for Figures 3-9.





Figure 19. Total Phosphorus content of P. boryanum as a function of culture age. Cultures were maintained at 37°C.



polyphosphate present should be at its lowest level (refer to Aspects of Phosphate Metabolism Section of Literature Review). Since the length of time necessary to deplete the cell of any phosphorus reserves was not known, a series of experiments were set up to determine the effects of phosphate starvation on rates of phosphate uptake. Cells taken from logarithmic phase were inoculated into phosphate-free medium for 3, 5, 11, and 12 days. The cells were then reinoculated into medium containing a known amount of phosphate, and the cells were then analyzed for total phosphorus content. These results are presented in Figure 20.

Phosphate uptake is represented by increase in total cell phosphorus for these experiments. Virtually no uptake occurred after 3 days of starvation (see legend to Figure 20 for environmental parameters). After 5 days of starvation, total cell phosphorus at the end of the uptake period increased from approximately 5 to 50 ug phosphorus per mg dry weight algae. Prolonging the starvation period to 12 days had no significant effect on the uptake rate. Starvation of the algae for 11 days and subsequent inoculation into a lower phosphate concentration (10 mg PO_4 per liter as opposed to 92.50 mg PO_4 per liter as above) resulted in uptake, but with reduced magnitude. Thus, 5 days of incubation in phosphate-free medium at 25°C, 500 ft-candles of illumination, and a 12 hour alternating day/night cycle was chosen to be the minimum time length necessary to induce phosphate starvation.

Determination of Uptake Parameters


After the determination of starvation conditions, it was

Figure 20. Semi-logarithmic plot of increase in total cell phosphorus as a function of increasing the length of time in phosphate-free medium (starvation). All uptakes were at 25°C, and 500 ft-candles of illumination. Closed squares represent 12 days of starvation and subsequent inoculation into medium containing 92.51 mg PO₄ / liter.

Closed Circles represent 5 days of starvation and uptake from 92.51 mg PO₄ / liter.

Open Squares represent 11 days of starvation and uptake from 10 mg PO₄ / liter.

Closed Triangles represent 3 days of starvation and uptake from 10 mg PO₄ / liter.



necessary to select parameters to monitor for the uptake of phosphate. The first two physical parameters considered were the dry weight of the algae and loss of phosphate from the medium (refer to Figure 22). Orthophosphate in the medium was essentially depleted at the end of the 4 hours of uptake. During this same period of time, the dry weight of the algae increased. Studies correlating cell number and dry weight analysis of normal, starved, and uptake cells indicate that the dry weight of the cells decreases during starvation and subsequently increases during rapid uptake, but not to the levels of normally grown cells (Table 3).

Table 3. Changes in Cell Size and Weight During Starvation and

	Cell Length	Dry Weight (mg)	Average Wt. mg/cell	Cells/mg dry weight
Normal	4.01 \pm 1.93 μ m	330	1.98 x 10 ⁻⁷	5.5 x 10 ⁶
Starved	2.87 \pm 0.73 μ m	60	2.20 x 10 ⁻⁸	
Uptake	2.80 \pm 0.69 μ m	200	7.48 x 10 ⁻⁸	

The average cell size decreased during starvation and uptake, indicating unusual and/or rapid cell division. During the starvation period, the average weight of the cell decreased by an order of magnitude. This value increased during uptake, but not to the value of a normal cell. These values were obtained by counting 300 filaments averaging 8 cells per filament, and taking dry weight measurements of the culture. For most cultures, 1 mg of dry weight corresponds to 5.5 x 10⁶ cells.

Figure 21. Increase in Total Cell Phosphorus and Cold TCA Extractable Phosphorus during Rapid Uptake at 25°C, 500 ft-candles of illumination, and 10 mg PO₄ / liter.

Figure 22. Increase in dry weight of the algae and depletion of medium phosphorus during rapid uptake as described for Figure 21.





Figure 21 demonstrates that during the 4 hour uptake period, the largest increase in total cell phosphorus and cold TCA extractable phosphorus occurs at 1 hour. The total cell phosphorus increases from about 8 to 12 ug P/mg dry weight and the cold TCA extractable phosphorus increases from 1 to 7 ug P/mg dry weight algae. This uptake then begins to level off, and is virtually complete at four hours. It was for these reasons that uptake was usually monitored for a four hour period.

Physical Parameters Affecting Phosphate Uptake Rates

A. Temperature

Rapid uptake of orthophosphate by the algae was monitored at three different temperatures, 4, 25, and 37°C. The 25° and 37°C uptakes were carried out on algae acclimated to those respective temperatures as previously described. Since the algae did not grow at 4°C, they were grown at 25°, and chilled for 1 hour prior to inoculation into a phosphate-containing medium. These results are presented in Figure 23. This graph demonstrates that there was little uptake at 4°C, and a definite reduction in total cell phosphorus after 2 hours of incubation at this temperature. The uptake response at 37° was essentially the same as that of 25°C. This was unlike the growth rate response at 37°C, which was a doubling.

B. Light


For all rapid uptakes at different light intensities, the algae were incubated during the starvation period at 500 ft-candles of illumination and then incubated at different light intensities

Figure 23. Semi-logarithmic plot of the increase in total cell phosphorus vs. uptake time as a function of temperature.

Circles represent uptake at 4°C.

Triangles represent uptake at 25°C.

Squares represent uptake at 37°C.



23

for the uptake. Variations in light intensity were accomplished by placing the culture flasks at different levels from the light source in the growth chamber, or in the case of total darkness, by covering the flasks entirely with aluminum foil. Figure 24 shows the results of these experiments. In total darkness or at 100 ft-candles, there was little uptake of orthophosphate, as expressed by increase in total cell phosphorus. The total cell phosphorus increased when the algae were exposed to orthophosphate at 500, 1000, and 2000 ft-candles of illumination. This increase was directly related to the light intensity. The greatest increase in total cell phosphorus for all light intensities occurred within the first hour of uptake (Figure 24).


C. Effect of Phosphate Concentration on Uptake

Monitoring the phosphate depletion from the medium led to studies which were designed to test the maximum amount of phosphate that could be utilized by the algae. It was found that the algae could essentially deplete the medium of phosphorus if given limited amounts (Figures 22, 25, and 26). These uptakes were carried out in medium containing 10 mg PO_4 (3.26 mg P) per liter or less. By increasing the amount of orthophosphate available in the medium to 92.51 mg per liter, the algae removed more phosphate from the medium, and the total cell phosphorus of the algae also increased (Figure 27). These results are summarized in Table 4.

D. Effects of Inhibitors on Uptake

For all studies of inhibitors, the algae were pre-incubated with the inhibitor for one hour prior to the addition of phosphate.

Figure 24. Increase in Total Cell Phosphorus during rapid uptake as a function of light intensity. The graphs from top to bottom, represent 2000 (○), 1000 (□), 500 (■), 100 (▼), and 0 (●) ft-candles of illumination



24

Figures 25-27. Change in total cell phosphorus and phosphate depletion of the medium as a function of orthophosphate concentration in the medium. Total cell phosphorus is represented by open symbols, and phosphate depletion of the medium by closed symbols.

Figure 25. 6.58 mg PO₄ / liter culture medium

Figure 26. 10.13 mg PO₄ / liter culture medium

Figure 27. 92.51 mg PO₄ / liter culture medium




Table 4. Summary of phosphate depletion in the medium and increase in total cell phosphorus as a function of external orthophosphate depletion.

TABLE 4

MEDIUM PHOSPHATE [mg./l]	MEDIUM PHOSPHORUS [mg./l]		DRY WEIGHT [mg./l]	Δ MEDIUM PHOSPHORUS	LARGEST INCREASE IN TOTAL CELL PHOSPHORUS	PHOSPHORUS IN MEDIUM AFTER 4 HOURS OF UPTAKE
	<u>conc.</u>	<u>time</u>		DRY WEIGHT [mg. P / mg. dw]		
6.58	2.15	0	20	1.03×10^{-1}	9.36×10^{-2}	8.5×10^{-2}
	0.09	4				
10.13	3.30	0	80	3.7×10^{-2}	1.19×10^{-2}	3.3×10^{-1}
	0.33	4				
92.51	30.16	0	75	7.4×10^{-2}	2.73×10^{-2}	24.65
	24.65	4				

Figure 28 illustrates the results of these experiments. There was virtually no uptake when 1×10^{-3} sodium fluoride, 1×10^{-2} mercuric chloride, or 5×10^{-3} 2,4,dinitrophenol were used as inhibitors. Of the three tested, sodium fluoride was the least effective. The upper curve in Figure 28, an uptake with no inhibitors present, under otherwise identical conditions, is plotted as a reference curve.

Normal Levels of Phosphorus-Containing Compounds
in Plectonema boryanum

After preliminary data were obtained about the "overplus" phenomenon in P. boryanum, studies were conducted to determine the distribution of phosphorus in various extracts. Nucleic acids as determined in each of these fractions, and determined for whole cell digests was no greater than 10^{-6} mg P per mg dry weight in any case, which is considerably less than any fraction reported. It is for this reason that they are not included in the data.

P. boryanum was grown for 14 days to logarithmic phase under normal culture conditions in Modified Fitzgerald containing various concentrations of phosphate. The algae were then extracted as previously described, and the extracts were measured for total phosphorus concentration. These results are presented in Table 5. Average values for triplicate samples, varying no greater than 1 per cent transmittance, in three different extractions are summarized in this table. Residues from each successive extraction were also tested as a check on the extraction procedure.

It can be seen that the phosphate levels for the total cell

Figure 28. Effect of metabolic inhibitors on the rapid uptake process. Semi-logarithmic Plot of increase in total cell phosphorus versus time. Variation of phosphate levels at time 0 can be explained by the standard deviations encountered during sampling.

Open Circles represent an uptake in the absence of inhibitors.

Closed Circles represent 1×10^{-3} M sodium fluoride

Closed Triangles represent 1×10^{-2} M mercuric chloride

Closed Squares represent 5×10^{-3} M 2,4-dinitrophenol




Table 5. Phosphorus Content of Cell Fractions as a function of a phosphate concentration in the culture medium. Values reported are mg P / mg dry weight.

TABLE 5

Phosphate in Medium	Phosphorus Content of Cell Fractions						
	TOTAL CELL DIGEST	COLD TCA EXTRACTION	RESIDUE FROM COLD TCA	LIPID EXTRACTION	LIPID FREE RESIDUE	HOT TCA EXTRACTION	RESIDUE
1 mg.	3.01×10^{-3} $\pm 0.38 \times 10^{-3}$	1.72×10^{-3} $\pm 0.36 \times 10^{-3}$	1.97×10^{-3} $\pm 0.32 \times 10^{-3}$	1.20×10^{-4} $\pm 0.33 \times 10^{-4}$	1.44×10^{-3} $\pm 0.25 \times 10^{-3}$	1.56×10^{-3} $\pm 0.18 \times 10^{-3}$	1.13×10^{-3}
10 mg.	8.25×10^{-3} $\pm 3.0 \times 10^{-3}$	5.51×10^{-3} $\pm 2.33 \times 10^{-3}$	5.85×10^{-3} $\pm 1.40 \times 10^{-3}$	7.54×10^{-4} $\pm 1.67 \times 10^{-4}$	6.71×10^{-3} $\pm 2.16 \times 10^{-3}$	4.49×10^{-3} $\pm 2.08 \times 10^{-3}$	3.23×10^{-3} $\pm 1.07 \times 10^{-3}$
100 mg.	3.28×10^{-2} $\pm 1.0 \times 10^{-2}$	2.79×10^{-2} $\pm 2.31 \times 10^{-2}$	7.58×10^{-3} $\pm 1.31 \times 10^{-3}$	3.37×10^{-4}	6.54×10^{-3} $\pm 2.06 \times 10^{-3}$	5.67×10^{-3} $\pm 0.72 \times 10^{-3}$	1.89×10^{-3} $\pm 1.34 \times 10^{-3}$
1000 mg.	3.58×10^{-2} $\pm 1.46 \times 10^{-2}$	2.00×10^{-2} $\pm 1.85 \times 10^{-2}$	4.80×10^{-3} $\pm 3.08 \times 10^{-3}$	2.30×10^{-4}	2.53×10^{-3}	1.95×10^{-3}	1.27×10^{-3}

digest, cold TCA extraction, and hot TCA extraction increase with increasing external orthophosphate concentration. This effect is not seen with the lipid extractions or residue. The phosphorus levels in these fractions were highest when the algae were grown in 10 mg PO_4 per liter. Since the algae were normally cultured in medium containing between 8 and 10 mg PO_4 per liter, the values listed for these concentrations in Table 5 were used as the normal phosphorus levels in these fractions.

Phosphorus Distributions During Starvation and Rapid Uptake

Table 6 summarizes the changes in phosphorus content of the various extracts during five days of starvation and four hours of rapid uptake. These results are also presented in Figures 29 - 33. It was not possible chemically to extract and analyze hourly all fractions during rapid uptake from one sample due to the length of time required for one extraction. The time required for one total extraction is approximately four hours. Consequently, 0, 2, and 4 hours of uptake were monitored in all phosphorus containing fractions whereas 1 and 3 hours were monitored for total cell phosphorus.

A. Total Cell Digest

Figure 29 demonstrates the changes in phosphorus content of the algae during starvation and uptake. The values of these fractions are also presented in numerical form in Table 6. During the 5 day starvation period, the total phosphorus content varied along with the dry weight of the algae. Five days of starvation


Table 6. Phosphorus Content of Cell Fractions during 5 days of phosphate starvation and 4 hours of phosphate uptake. Sample dry weights for starved cells are presented to demonstrate the fluctuations in dry weight that occur during starvation. The increase in dry weight in a representative uptake can be seen in Figure 22. Values reported are mg P / mg dry weight.

growth conditions	TOTAL PHOSPHORUS IN CELL FRACTIONS							
	Dry Wt.	Total Cell Digest	Cold TCA	Cold TCA Residue	Lipid Extract	Lipid Free Residue	Hot TCA Extract	Residue
Normal Cells 8-10mg.		8.25×10^{-3} $\pm 3.00 \times 10^{-3}$	5.51×10^{-3} $\pm 2.33 \times 10^{-3}$	5.85×10^{-3} $\pm 1.40 \times 10^{-3}$	7.54×10^{-4} $\pm 1.18 \times 10^{-4}$	6.71×10^{-3} $\pm 2.16 \times 10^{-3}$	4.49×10^{-3} $\pm 2.08 \times 10^{-3}$	3.23×10^{-3} $\pm 1.07 \times 10^{-3}$
1 Day Starved	1200	5.35×10^{-3}	2.52×10^{-3}		1.09×10^{-4}		2.52×10^{-3}	1.75×10^{-3}
2 Day Starved	2400	2.48×10^{-3}	1.17×10^{-3}		7.0×10^{-5}		1.29×10^{-3}	8.7×10^{-4}
3 Day Starved	1400	5.38×10^{-3}	1.97×10^{-3}		9.3×10^{-4}		1.91×10^{-3}	3.12×10^{-3}
4 Day Starved	2000	2.98×10^{-3}	1.27×10^{-3}		1.4×10^{-4}		1.21×10^{-3}	1.72×10^{-3}
5 Day Starved		4.19×10^{-3} $\pm 0.92 \times 10^{-3}$	2.35×10^{-3} $\pm 1.08 \times 10^{-3}$	5.73×10^{-3}	1.03×10^{-4}	5.1×10^{-3}	2.68×10^{-3} $\pm 1.56 \times 10^{-3}$	3.3×10^{-3} $\pm 2.12 \times 10^{-3}$
1 Hour Uptake		4.43×10^{-2} $\pm 1.86 \times 10^{-2}$	7.02×10^{-3}					
2 Hour Uptake		4.78×10^{-2} $\pm 1.96 \times 10^{-2}$	7.70×10^{-3}		3.5×10^{-3}			8.6×10^{-3}
3 Hour Uptake		4.69×10^{-2} $\pm 1.75 \times 10^{-2}$	7.20×10^{-3}					
4 Hour Uptake		7.83×10^{-2} $\pm 1.52 \times 10^{-2}$	2.35×10^{-2} $\pm 0.77 \times 10^{-2}$	4.40×10^{-3}	5.6×10^{-3}		3.88×10^{-2}	9.0×10^{-3}

102

TABLE 6

Figure 29. Semi-logarithmic graphic presentation of the change in Total Cell Phosphorus during starvation and rapid uptake. The standard error of the samples is presented in this figure and in all subsequent figures as open circles connected by lines. Standard errors are plotted where four or more samples were analyzed. All other data represents the mean of three extractions.



led to a reduction of the total phosphorus content of the algae by approximately one-half. Upon the addition of phosphate, this value increased by an order of magnitude at one hour and showed an additional small increase at four hours.

B. Cold TCA Extractable Phosphorus

The values for this fraction follow a similar pattern for the starvation period (Figure 30 and Table 6). Again, the phosphorus content of the cold TCA extract was reduced by about one-half at the end of the 5 day starvation period. However, the most substantial increase was found at four hours of uptake, when the value was one order of magnitude higher than that of the value for 5 days of starvation.

C. Lipid Extraction

The phosphorus content of the lipid fraction was reduced at the end of the five day starvation period by 81 per cent (Figure 31 and Table 6). At the end of the 4 hour uptake period, this value did not return to the levels found in normal cells, indicating that the phosphate lost during starvation was not recovered in this fraction within 4 hours.

D. Hot TCA Extractable Phosphorus

The greatest increase in phosphorus content of any cell fraction during the uptake period was found in the hot TCA extract (Figure 32 and Table 6). Five days of starvation led to a reduction in this fraction by approximately 40 per cent. At the end of the 4 hour uptake period, the increase in phosphorus content was greater than an order of magnitude, the value increasing from 2.68 to 38.8 ug P

**Figure 30. Semi-logarithmic graphic presentation of the change
in cold TCA extractable phosphorus during starvation and rapid uptake.**




Figure 31. Semi-logarithmic graphic presentation of the change in phosphorus content of lipid fraction during starvation and rapid uptake.





Figure 32. Semi-logarithmic graphic presentation of the change in hot TCA extractable phosphorus during starvation and rapid uptake. Hourly sampling of the hot TCA extract was difficult, and the results obtained were questionable. Therefore, only the sample taken on the fourth hour was plotted.

Mg. phosphorus / mg. dry weight



per mg dry weight algae.

It is also interesting to note that although in normal cells the amount of phosphorus present in the hot TCA extract was less than that in the cold extract, the values at 4 hours of uptake were reversed. There was more phosphorus present in the hot TCA extract. During the five day starvation period, these values were approximately the same.

E. Residue from the Threefold Extraction


The residue from the threefold extraction as previously described showed the greatest variation in phosphorus content during the starvation period (Figure 33 and Table 6). However, the phosphorus content of this fraction was essentially the same for a 5 day starved culture and a normal culture. The increase after 4 hours of uptake was also not as large as the other extractions. This fraction probably represents all phosphorus compounds in the cell other than some nucleic acids, short and long chain polyphosphates, orthophosphate, and phospholipids.

Hydrolysis of Condensed Phosphates

Hydrolysis of the condensed phosphate, sodium tripolyphosphate, was tested both with and without the experimental organism. Modified Fitzgerald's medium was made with three different concentrations of the condensed phosphates, 4.59, 153.7, and 1204 mg PO_4 per liter culture medium. The tubes were incubated under normal culture conditions as previously described. These results are presented in Figures 34 - 36.

Figure 34 demonstrates that at a concentration of 4.59 mg PO_4

Figure 33. Semi-logarithmic graphic presentation of the change in phosphorus content of the residue from the threefold extraction during starvation and rapid uptake.




Figures 34-36. Hydrolysis of sodium tripolyphosphate in Modified Fitzgerald medium both in the presence and absence of P. boryanum. Circles represent hydrolysis without cells, and squares represent hydrolysis with cells present. Hydrolysis is indicated by an increase in orthophosphate present in the medium.

Figure 34. 4.59 mg PO₄ / liter culture medium

Figure 35. 153.7 mg PO₄ / liter culture medium

Figure 36. 1204 mg PO₄ / liter culture medium



per liter, the condensed phosphate apparently hydrolyzed at a greater rate with the alga present in the medium. At a concentration of 153.7 mg PO_4 per liter (Figure 35), the condensed phosphate hydrolyzed more rapidly in the presence of the alga. However, this value began to decrease on the tenth day of culture, and did not rise again. Hydrolysis in all cases was determined by an increase in orthophosphate concentration in the medium. Figure 36 illustrates the rapid hydrolysis of a considerable amount of condensed phosphate on the first day of culture, and a subsequent steady increase in the amount of orthophosphate present in medium. At the higher concentrations of condensed phosphate (Figures 35 and 36), hydrolysis in the absence of algae proceeded at a constant rate.


Rapid Uptake of Condensed Phosphates

Sodium tripolyphosphate was substituted for orthophosphate in a normal starvation and rapid uptake experiment. Total cell fractionations of phosphorus-containing compounds were assayed. These results are presented in Figure 37. There was no substantial increase in any cell fraction after four hours of uptake. This is in contrast to an uptake with orthophosphate as previously discussed. Refer again to Table 6 and Figures 29-33.

Cell Ultrastructure Under Normal Culture Conditions

Cells of P. boryanum grown under normal culture conditions have an appearance which is similar to that of other blue-green algae. Cellular inclusions such as polyhedral bodies, areas of DNA, ribosomes, thylakoids, lipid droplets, cyanophycean granules, and a few small

Figure 37. Increase in phosphorus-containing fractions during starvation and rapid uptake from 10 mg PO_4 (as sodium tripolyphosphate) per liter. The left bar of each pair represents starved cells, and the right bar represents 4 hours of uptake. The uptake with condensed phosphates was only repeated twice. Consequently the values reported represent the mean.



polyphosphate bodies are found in the cell with regularity (Figures 38 and 39). A four-layered cell wall, as described in most blue-green algae is also present (Figure 39). Some intrathylakoidal vacuolization is present, with the thylakoids usually arranged at the periphery of the cell (Figure 38). Cell division occurs by transverse binary fission. A newly formed septum can be seen in Figure 38.

Changes in Ultrastructure Associated with Several
Phosphate Concentrations

A. 100 mg PO₄ per liter

P. boryanum grown in culture medium containing 100 mg PO₄ per liter had a number of alterations in cell structure. Figure 40 demonstrates these changes. Generally, there is a marked increase in cell death and cell lysis, as evidenced by the debris in the sectioned pellet. The cells which appear to be dead have cytoplasm which is devoid of ribosomes. The thylakoids are also fewer in number and are rather prominent in the cell. Occasionally, large polyphosphate bodies, or areas where polyphosphate bodies have fallen out, can be seen. Cells which are normal in appearance, as previously described, can also be found.

B. 1000 mg PO₄ per liter

Changes that occurred in culture at 100 mg per liter were even more pronounced when the algae were grown in medium containing 1000 mg PO₄ per liter (Figures 41-43). Figure 43 demonstrates some expansion of thylakoids (IT), and a large hole in the center of the cell, presumably an area where a polyphosphate body has fallen out during

Key to Figure Legends of Electron Micrographs

- C = Cyanophycean Granule**
- D = DNA**
- EL = Electron Lucent Area**
- IT = Intrathylakoidal Spaces**
- L = Lipid**
- P = Polyphosphate Body**
- Pb = Polyhedral Body**
- PM = Plasma Membrane**
- R = Ribosomes**
- S = Septum**
- Sh = Sheath**
- T = Thylakoids**

Figure 38. Cells of Plectonema boryanum grown for 14 days in 10 mg PO_4 / liter at 500 ft-candles of illumination with an alternating 12 hour day/night cycle. Note the relatively small intrathylakoidal spaces, and inclusions normally present in blue-green algae such as lipid droplets (L), polyhedral bodies (Pb), and areas of DNA (D). Cell division which will result in two equal daughter cells can also be seen (S).

Figure 39. P. boryanum grown under normal conditions described above and shown at a higher magnification. Note the arrangement of the cell wall outside the plasma membrane (PM). Layer 2 is the mucopolymer-containing layer.





Figure 40. Low power picture of a sectioned pellet of P. boryanum grown in Modified Fitzgerald's Medium for 14 days containing 100 mg PO_4 / liter at 500 ft-candles of illumination. Note the cellular debris in the pellet, and the appearance of several types of cells. The filament in the upper left corner of the picture can be considered as "normal", resembling cells grown under the conditions listed for Figure 38. Other cell types are also evident.




Figures 41-43. Cells of Plectonema boryanum grown for 14 days in medium containing 1000 mg PO_4 / liter at 500 ft-candles of illumination. Note the cell types present, as in the culture containing 100 mg PO_4 / liter.

Figure 41. Portions of cells showing some expansion of intrathylakoidal spaces (IT), thylakoids (T) at the periphery of the cells, and DNA (D) and polyhedral bodies (Pb) surrounding a space where a polyphosphate body has fallen out.

Figure 42. A cell, in a stage of degradation, which is devoid of ribosomes. Note the small lipid droplets (L) throughout the cell, and the prominent thylakoids (T).

Figure 43. Portion of a filament showing numerous polyphosphate bodies (P) and enlarged ribosomes (R).



sectioning. In the more normal appearing cells, the thylakoids are still located at the periphery, and the DNA is located in the center of the cell. Figure 42 shows in greater detail a cell which appears to be dead and is devoid of ribosomes. The frequency of encountering such cell types increases with greater phosphate concentrations. Figure 43 shows a portion of a filament containing several polyphosphate bodies which have sublimated under the electron beam or have been partially lost during the section preparation procedures. The ribosomes also have an unusual appearance, being much larger than in cells grown in normal culture medium.

C. 1 mg PO₄ per liter

Cells grown in culture medium containing 1 mg PO₄ per liter have approximately one-tenth of the normal phosphate concentration available to them. This particular culture condition also leads to changes in cell architecture (Figures 44-47), but these changes are unlike those of higher phosphate concentrations. The most pronounced changes that occur are the expansion of intrathylakoidal spaces (IT) and the development of areas of medium electron density (arrows in Figures 44 and 45). In these figures, the appearance of the cytoplasm and some of the ribosomes is normal in all other respects. Some ribosomes (Figure 47) are unusually large. Polyhedral bodies and areas of DNA are also present. Elongate cells were frequently encountered (Figure 47). These cells often attained lengths of 6 - 9 μ m, whereas a normal cell was about 3 μ m. They frequently have abnormally large ribosomes. Refer again to Figures 46 and 47.



Figures 44-47. Cells of Plectonema boryanum grown in culture medium for 14 days containing 1 mg PO₄ / liter at 500 ft-candles of illumination and a 12 hour alternating day/night cycle.

Figure 44. Cell showing expansion of intrathylakoidal spaces (IT) and the presence of an area of medium electron density (arrow). DNA (D) and polyhedral bodies (Pb) are located in the center of the cell.

Figure 45. Cell showing a greater degree of expansion of intrathylakoidal spaces (IT). An immature polyphosphate body (arrow) can also be seen.

Figure 46. A second cell type found in phosphate-limited cultures. These cells are characterized by unusually large ribosomes (R).

Figure 47. Portion of an elongate cell of P. boryanum. The cell contains many small lipid droplets (L) and ribosomes (R).



Cells Grown in the Absence of Phosphate


Cells grown in the absence of phosphate, previously described as a 5 day starvation period, had many structural similarities to cells grown in 1 mg PO_4 per liter for 14 days. The most obvious change is the increasing frequency with which these changes were found with progression further into the starvation period. Figures 48 - 54 illustrate these changes.

The cell division process appears to be abnormal in cases of phosphate limitation. This leads to two unusual conditions: 1. the presence of elongate cells (Figure 48) and 2. the presence of unequal cell divisions (Figure 49). The unequal cell divisions often resulted in cells which were no greater than 0.4 μ m in length. A great number of divisions have been observed wherein the newly forming septa are not opposite (Figure 49). This results in a bend of the filament at that point.



Another prominent change that occurs during phosphate starvation was the expansion of intrathylakoidal spaces (Figures 48 - 52). The degree of expansion seems to be related to the length of time in phosphate-free medium. Relatively small intrathylakoidal spaces (Figures 48 and 49) enlarge to areas such as those seen in Figures 50 and 51. During the starvation period and most evident at the fifth day of phosphate starvation were the development of areas of medium electron density (EL) such as that shown in Figure 52. On a relative basis, the number of cells with no thylakoid expansion (Figure 52) was less. Such cells most often contain ribosomes which appear larger than normal.

Figure 48. Elongate cell of P. boryanum grown in phosphate-free medium for 2 days. Note several small polyphosphate bodies (P), and the increased vacuolization. DNA (D), polyhedral bodies (Pb), and lipid droplets (L) are also present. The cell is approximately 6 um in length.

Figure 49. Portion of a filament of P. boryanum starved of phosphate for 3 days. Note the unequal cell divisions and septa (arrows, S) which are not opposite. Some sheath (Sh) is also present.



Figures 50-52. Cells of P. boryanum grown in phosphate-free medium for 5 days. Note the large expansion of intrathylakoidal spaces (IT), and the development of an electron-lucent area (EL) in the center of the cell. Unusual looking ribosomes (R) are also present in these cells (Figure 52).



In cells cultured in all phosphate concentrations, little (as in Figures 40, 49, 50 and 54) or no sheath (for example, Figures 38, 39, 40, and 41) was present.

Another structural feature of phosphate starvation was the presence of large lipid-like inclusions in the cell (Figures 53 and 54). These were in contrast to the smaller lipid droplets appearing in Figure 48. These larger lipid-like inclusions appeared in both normal (Figure 53) and degenerating cells (Figure 54).

Changes in Ultrastructure During Rapid Uptake

The rapid uptake process, or "overplus phenomenon", was also quite distinct at the ultrastructural level. The most notable changes were the large areas of medium electron density and the increase in number and size of polyphosphate bodies. The location and association of polyphosphate bodies with other cellular inclusions were also novel.

Phosphate limitation, as previously discussed, resulted in the expansion of intrathylakoidal space and the development of areas of medium electron density (EL). This area expanded even further during the uptake process, and was in fact, quite characteristic of cells in this metabolic condition (Figures 55 - 63). The developing large polyphosphate bodies were usually located within these areas (Figures 59 - 62). Polyphosphate bodies are quite dense and usually chip out or fall out during sectioning, leaving characteristic holes which frequently have some polyphosphate remaining in the epoxy (Figures 55, 57, 58, and 63). Oftentimes, one can observe strands which may be DNA seemingly attached to, or associated with, the

Figures 53-54. Two cell types of P. boryanum from a culture starved of phosphate for 3 days. Note the presence of large lipid-like inclusions (L). Cyanophycean granules (C) are also present.





Figure 55. Cell of P. boryanum at 2 hours of uptake. Note the large electron-lucent area (EL) in the center of the cell, and some polyphosphate remaining in the section. Strands of DNA-like material are also present in the electron-lucent area. All cells in the following figures were taken from cultures which had been starved of phosphate for 5 days and subsequently inoculated into medium containing 10 mg PO₄ / liter. Samples were withdrawn and fixed hourly.

Figure 56. Large electron-lucent area in an uptake cell with DNA (D) at the periphery.








Figure 57. Electron-lucent area (EL) containing holes where polyphosphate bodies have seemingly been lost, in an uptake cell with unusually large ribosomes (R). A polyhedral body (Pb) can also be seen.

Figure 58. Large electron-lucent areas (EL) in two cells, with numerous polyphosphate bodies (P) in various stages of development. Note the presence of a developing polyphosphate body (P) in a polyhedral body (Pb).



Figures 59-62. Cells of P. boryanum at 4 hours of uptake showing large polyphosphate bodies (P) which have not fallen out in sectioning. Note the "smearing" of polyphosphate bodies (P) in the direction of sectioning.



polyphosphate body (Figures 55 and 57). The electron lucent area was always located in the center of the cell, with the DNA either dispersed in this area (Figures 55 and 57) or at the periphery of this area (most easily observed in Figures 56 and 61). The appearance of the ribosomes was as in the phosphate limited cells (Figures 57 and 58), with normal and enlarged ribosomes present in 2 cell types.

Cells may accumulate one large polyphosphate body (Figures 59 - 63) or several smaller polyphosphate bodies (Figures 57-59), which can be seen in thin sections. One can observe occasionally polyphosphate bodies located in polyhedral bodies (Figure 58). In addition to the large mature bodies, smaller bodies in various stages of development are also found in the cells (Figure 58). Figures 59 - 63 show large polyphosphate bodies which have not chipped out in sectioning. The bodies are extremely electron dense, and often smear in the direction of sectioning (Figures 61 and 62). One can observe, but with lesser frequency, entire filaments that contained polyphosphate bodies (Figure 63). This is an orientation problem associated with sectioning the algae.




Light Microscopy

Cells stained by the lead sulfide precipitation method after 5 days of phosphate starvation contain a few small polyphosphate bodies (Figure 64). Cells which have undergone the uptake of phosphate for 4 hours after the starvation period usually contain one apparent large polyphosphate body located in the center of the cell (Figure 65).

Figure 63. Portion of a filament of P. boryanum at 4 hours of uptake showing electron-lucent areas and remnants of polyphosphate bodies.

Figure 64. Light micrograph of a filament of P. boryanum starved of phosphate for 5 days and stained for polyphosphates.

Figure 65. Light micrograph of a filament of P. boryanum stained for polyphosphates at 4 hours of phosphate uptake. Note the presence of large polyphosphate bodies in all cells.



X-ray Energy Dispersive Analysis

X-ray energy dispersive analysis was performed on cultures of P. boryanum starved of phosphate for 5 days and inoculated into medium containing 10 mg of phosphate per liter. Thick sections (approximately 0.5 μm) were then examined as previously described. These results are presented in Figures 66-75. The fundamental principles underlying x-ray energy dispersive analysis are as follows: X-rays, characteristic of given elements, are produced by means of charged particles (e.g. electrons) bombarding that given element. The energies, as well as wavelengths, of the x-rays emitted upon bombardment are characteristic of each element due to transitions of electrons between energy levels and due to the charge of the nucleus. It is possible to detect both qualitatively and semi-quantitatively the presence of an element due to these x-ray energy emissions.

Figures 66-68 demonstrate the results of energy dispersive analysis of cells of P. boryanum fixed in glutaraldehyde and embedded in Epon. Figure 66 is an analysis of a portion of cytoplasm with no discernible polyphosphate bodies. Background emission, and a major and minor peak are evident. The minor peak corresponds to an energy of 2.136 keV, representing a $K\beta_1$ emission of phosphorus. The major peak is an energy of 2.61 keV, corresponding to a $K\alpha_1$ emission of chlorine. Chlorine is a common element encountered in epoxy resins.




Figure 67 is an x-ray energy dispersive analysis of a polyphosphate body, comparable in size and density to those in Figures 59-62. Analysis showed that the polyphosphate body consisted of phosphorus, character-

Figures 66-68. X-ray energy dispersive analysis of P. boryanum starved of phosphate for 5 days and inoculated into medium containing 10 mg PO₄ / liter. Samples are fixed in glutaraldehyde and embedded in Epon.

Figure 66. Analysis of cytoplasm containing no polyphosphate bodies. Minor peak: Phosphorus (2.136keV). Major peak: Chlorine (2.62keV).

Figure 67. Analysis of a dense polyphosphate body. Major peaks: Copper (1.10 and 8.04keV), phosphorus (2.01keV), chlorine (2.62keV), calcium (3.68keV).

Figure 68. Analysis of a developing polyphosphate body. Major peaks: copper (1.10 and 8.04keV), phosphorus (2.01keV), chlorine (2.62keV), calcium (3.68keV).



ized by $K\alpha_1$ and $K\alpha_2$ emissions of 2.013 and 2.012keV, and calcium, characterized by $K\alpha_1$ and $K\alpha_2$ emissions of 3.691 and 3.687keV. Other peaks which are present and minor are those of chlorine (2.62keV) and copper (8.04keV and 1.10keV). The 1.10keV emission of copper is a critical absorbance value, and 8.05keV is that of $K\alpha_1$ emission. The peaks for emission from copper are present because the sections were mounted on copper grids. Thus, the only elements present in any great amount are phosphorus and calcium. Identification of elemental analysis in the following descriptions will be limited to the element identified, and the energy characteristic of that element. No further references will be made to the specific emissions.

Figure 68 is an analysis of a polyphosphate body which has an internal area that is light and somewhat porous. The periphery of the body is dense. This type of polyphosphate body has been previously described as a developing stage (Jensen, 1969). The major elements present are phosphorus (2.01keV) and calcium (3.68keV). Other components are chlorine (2.62keV) and copper (8.04 and 1.10keV).

Figures 69-71 are energy dispersive analyses of samples fixed in osmium and embedded in Epon. Samples were first analyzed after osmium fixation because the fixative employed usually gives better preservation than glutaraldehyde, or glutaraldehyde - osmium. However, difficulties were encountered in elemental analysis of phosphorus when osmium was present. This was due to an $M\beta$ emission of 1.978keV from osmium. One frequently encounters a shoulder on the phosphorus peak if osmium is employed as a fixative.

Figures 69-71. X-ray energy dispersive analysis of cultures of P. boryanum starved of phosphate for 5 days and inoculated into medium containing 10 mg PO_4 /liter. Samples were fixed in osmium and embedded in Epon.

Figure 69. Analysis of epoxy containing no cells. Major peak: Copper (8.05keV).

Figure 70. Analysis of cytoplasm containing no discernible polyphosphate bodies. Major peaks: Copper (1.10 and 8.04keV), osmium (1.97keV), phosphorus (2.01keV), and calcium (3.68keV).

Figure 71. Analysis of a dense polyphosphate body. Major peaks: Osmium (1.97keV), phosphorus (2.01keV), calcium (3.68keV), and copper (8.04keV).






Figure 69 is an analysis of the epoxy embedding resin. Background emission is present, as well as a major peak corresponding to copper (8.05keV).

Figure 70 is an analysis of cytoplasm, containing no discernible polyphosphate bodies. Some phosphorus and calcium are present, (2.01 and 3.68keV) as well as chlorine (2.62keV) and copper (1.10 and 8.05keV). It is difficult to ascertain the relative amount of phosphorus due to the interference by osmium present in the sample. Figure 71 illustrates quite clearly the shoulder on the phosphorus peak, contributed by the osmium. This was an analysis of a dense polyphosphate body. Other major components are calcium (3.68keV), chlorine (2.62keV) and copper (8.05keV). The height and number of copper emission peaks encountered are a function of the distance of the grid bar from the area being analyzed.




Figures 72-74 are analyses of samples embedded in Durcupan, a water-soluble embedding medium. Durcupan was employed to determine if any major components were extracted during the alcohol and propylene oxide dehydration series normally used with epoxy embedding. Figure 72 is a reference analysis of the embedding medium with no cells present. The only major peaks encountered are those of copper (8.05keV) and osmium (8.90keV). Figure 73 is an analysis of cytoplasm. Major peaks which are identifiable are those of osmium (1.98keV), chlorine (2.62keV), copper (8.04keV), and osmium (8.90keV). Figure 74 is an analysis of a dense polyphosphate body. In addition to the peaks identified above, phosphorus (2.01keV) and calcium (3.68keV) are present. Thus it appears that the major components of the polyphosphate

Figures 72-74. X-ray energy dispersive analysis of a culture of Plectonema boryanum starved of phosphate for 5 days and inoculated into medium containing 10 mg PO_4 /liter. Samples were fixed in osmium and embedded in Durcupan.

Figure 72. Analysis of Durcupan embedding medium. Major peaks: Copper (8.04keV) and Osmium (8.90keV).

Figure 73. Analysis of cytoplasm containing no discernible polyphosphate bodies. Major peaks: Osmium (1.98keV and 8.90keV), chlorine (2.62keV), and copper (8.05keV).

Figure 74. Analysis of a dense polyphosphate body. Major peaks: Osmium (1.98 and 8.90keV), phosphorus (2.01keV), calcium (3.68keV) and copper (8.05keV).



bodies examined which are identifiable by this method of analysis
are phosphorus and calcium.

DISCUSSION

Growth Curves, Phosphate Depletion, and Minimum Cellular Phosphate Levels

The growth curves of Plectonema boryanum and Oscillatoria tenuis all demonstrate an increase in dry weight on the first day after transfer into fresh medium, regardless of phosphate concentration in the medium. The increase in dry weight is also accompanied by a corresponding decrease in phosphate concentration of the external medium (Figures 3 - 16). This increase in dry weight and depletion of phosphate from the external medium is probably due to an uptake phenomenon. Transfers of cultures were routinely made from cultures which were at least 2 months old. Analysis of the total phosphate content of cells as a function of culture age revealed that at this age, cultures had already been reduced in phosphorus content to 4 ugP per mg dry weight algae (Figure 18). Studies correlating dry weight analysis to cell number revealed that for all cultures, 1 mg of dry weight algae corresponded to 5.5×10^6 cells. Thus, the total phosphorus content of a 2 month old culture would equal 0.73 ug P per 10^6 cells.

Similar analysis of a 14 day old (normal), 5 day starved, and 4 hour rapid uptake cultures yield the following information:

Normal cultures - 1.5 ug P / 10^6 cells
Starved cultures - 0.76 ug P / 10^6 cells
Uptake cultures - 14.2 ug P / 10^6 cells

It can be seen that the total phosphorus content of a two month old culture is slightly, but not significantly, lower than that of a culture starved for 5 days. Normal culturing of the algae also leads to depletion of phosphorus reserves, but over a much longer period of time. Thus, transfer of these phosphate depleted cells, in reality, sets the stage for a rapid uptake of orthophosphate.

These results for internal phosphorus concentration are similar to those reported for other organisms. Mackereth (1953) reported that the limiting phosphorus concentration of Asterionella formosa was about $0.06 \text{ ug P} / 10^6 \text{ cells}$. This diatom could also store phosphorus in concentrations as high as $7 \text{ ug P} / 10^6 \text{ cells}$. The marine diatom Phaeodactylum tricornutum was found by Kuenzler and Ketchum (1962) to contain phosphorus concentrations of $2.1 \text{ ug P} / 10^6 \text{ cells}$ at its highest levels, and $0.06 \text{ ug P} / 10^6 \text{ cells}$ at its lowest levels. Kholy (1956) reported that the minimum value below which growth did not occur in Chlorella was $0.1 \text{ ug P} / 10^6 \text{ cells}$. The maximum level of phosphorus in the cells was $1.5 \text{ ug P} / 10^6 \text{ cells}$. This value did not increase further when the starved algae were re-inoculated into higher phosphate concentrations.

The values reported in the literature are similar to those values obtained for Plectonema boryanum in this study. However, it would appear that P. boryanum contains more total phosphorus even at its lowest levels, especially when one considers that this alga is much smaller in size than the diatoms or green alga utilized in these previous studies.

Table 3 indicates that there is an appreciable increase in dry weight of the algae during the uptake process. This increase in dry weight is also common to routine culture transfer (Figures 3-16). These results also lend support to the occurrence of the uptake phenomenon associated with transfer of algae to fresh medium during routine culture.

Physical Parameters Affecting Uptake Rates

Light has been demonstrated to stimulate uptake of phosphate in many organisms (Talpasayi, 1962; Kanai and Simonis, 1968; Fitzgerald, 1970). Dark assimilation of phosphate has also been reported (Kanai, Miyachi, and Miyachi, 1963; Batterton and Van Baalen, 1968; and Overbeck, 1962). P. boryanum appears to be an organism which requires light for maximal uptake of phosphate (Figure 24). The increase in total cell phosphorus is directly related to the light intensity available. Higher light intensities during the uptake period result in higher cell phosphorus levels, up to $18.9 \text{ ug P} / 10^6$ cells at 2000 ft-candles. Preillumination of cultures of P. boryanum during the starvation period and subsequent inoculation into a phosphate-containing medium in the dark resulted in uptake, but at greatly reduced levels. These results are unlike those reported by Simonis and Urbach (1963) who reported dark uptake in preilluminated cultures of Ankistrodesmus braunii.

The effect of light intensity on the uptake process suggests several possibilities. Energy is required for phosphate uptake to occur. This energy can be supplied by photosynthesis, respiration, or a combination of both processes. If photosynthesis alone were

involved, one would expect results such as those presented in Figure 24. ATP produced during photophosphorylation would supply the energy required for uptake. Another possible source of energy would be the metabolism of products of photosynthesis. If this were true, then inhibition of metabolic pathways involved in utilization of photosynthetic products would result in either complete or partial inhibition of the uptake process. A third alternative is the metabolism of storage materials in the cell through respiration. In this case, one would expect increased uptake with increasing light intensity, and also phosphate uptake in the dark with the metabolism of storage materials.

Figure 24 demonstrates that increasing light intensity does accelerate phosphate uptake, but there is virtually no uptake in the dark. Figure 28 illustrates the effects of several metabolic inhibitors on the phosphate uptake process. These results indicate that phosphate uptake requires a supply of metabolic energy. If metabolism of the products of photosynthesis were not involved as an energy source, one would expect no effect exerted by these inhibitors. This does not appear to be the case. Increasing light intensity most likely increases the supply of sugars to be metabolized by the alga. Some dark uptake probably indicates that sugars resulting from photosynthesis during the starvation period were probably metabolized in the dark, but no further production was possible. It is also likely that there are no other storage forms of sugar present in the alga which could readily be available as an energy source.

It is a well known fact that the optimal temperature for photo-

synthesis is 25°C. Temperature effects on uptake (Figure 23) and growth rates (Figure 17) indicate that although elevated temperatures (25-37°C) increased the growth rate, they do not increase the rate of uptake.

It has been demonstrated many times that an energy source is required for phosphate uptake. Blum (1966) demonstrated that phosphate uptake by Euglena gracilis was inhibited by 2,4-dinitrophenol. Borst-Pauwels and Jager (1969) found that phosphate uptake in Saccharomyces cerevisiae was inhibited by both 0.1 mM 2,4-dinitrophenol, and 20 mM fluoride. Jungnickel (1970) found that 2,4-dinitrophenol reduced, but did not completely inhibit phosphate uptake in Candida utilis. Ullrich (1972) found that rates of polyphosphate formation were rather low when compared with the probable rates of ATP formation under various conditions of photophosphorylation in Ankistrodesmus braunii. The uptake rate of phosphate after a starvation period was higher than the rate of polyphosphate synthesis.

Phosphorus Distributions in Cellular Extracts

A. Logarithmic Cultures in Different Phosphate Concentrations

Table 5 summarizes the influence of external phosphate concentration on the distribution of phosphorus in various cell extracts. It can be seen that increasing phosphate concentrations, in general, result in higher phosphorus levels for all fractions. These results can also be correlated with the microscopy of the samples.

The ratios of acid-soluble (cold TCA extractable) phosphates to acid-insoluble (hot TCA extractable) phosphates seem to vary in many organisms. Aitchison and Butt (1973) reported that acid-insoluble

polyphosphates of Chlorella vulgaris were slightly higher during logarithmic growth than acid-soluble. Baker and Schmidt (1964) found that in Chlorella pyrenoidosa, the levels of acid-insoluble polyphosphate decreased immediately prior to and during nuclear division. Krishnan et al (1957) found that both acid-soluble and acid-insoluble polyphosphates were normally present in mycelia of Aspergillus niger. Soluble polyphosphate was present in low concentrations in new mycelia, but accumulated and reached a peak as the organism grew, after which it declined until the time of general autolysis. On the other hand, they found that insoluble polyphosphates were present in high concentrations in newly formed mycelia, and fell after about one week at these concentrations.

Plectonema boryanum, during logarithmic growth, contains more acid-soluble than acid-insoluble polyphosphates (Table 5). This is true for all phosphate concentrations tested. The standard deviations expressed in this table are a result of variability in different experiments, and not variability in the triplicate samples of a single total extraction. The increasing total cell phosphorus content is probably due to the uptake phenomenon occurring upon transfer to fresh medium. It has been demonstrated earlier that the algae, under optimal conditions, do not increase in total cell phosphorus to levels above 18.9 ug P per 10^6 cells. This maximum level occurred only at 2000 ft-candles of illumination and was not used routinely for uptake studies because the alga will not grow well at light intensities above 500 ft-candles.

In the case of growth in 1000 mg of phosphate per liter of

culture medium, the total phosphorus content of the cell digest represents 6.5 ug P per 10^6 cells, a value approximately four times higher than that of the cultures grown at 10 mg PO_4 per liter. If the uptake phenomenon did occur upon transfer, one would expect increased total cell phosphorus according to the available phosphorus in the medium, but probably reduced after 14 days of culture. The phosphorus content of the total cell digests of cultures grown in 100 and 1000 mg PO_4 per liter culture medium again also indicates that the algae are able to assimilate certain levels of phosphate, and beyond a certain value, no further increase in phosphate concentration in the medium results in increased cellular phosphate levels.

Electron microscopic examination of cultures grown at various phosphate concentrations also reveals that there is increased cell death and lysis when the algae are grown at high phosphate concentrations. This might indicate that the measurements of phosphorus concentrations in the cell fractions of cells grown in higher phosphate concentrations are more unreliable, since the number of viable cells is reduced.

B. Starved and "Overplus" Cultures

Table 6 and Figures 29-33 demonstrate the changes in all phosphorus-containing fractions that occur when the algae are cultured in phosphate-free medium and subsequently inoculated into a medium containing phosphate. Cultures starved of phosphate for a five day period generally decrease in phosphate levels in all cell extracts examined. After the initial reduction on days 1 and 2, there was an increase, though small, on days 3 and 5. This pattern

of changes in phosphorus content can be related to the fluctuations in dry weight which occur during the starvation period, and also to the microscopic examination of the cultures.

There is a loss in dry weight of the cultures on days 2 and 4 of starvation. The increases in phosphorus occur when there is a reduction in dry weight. Examination of the cultures during this period reveals that there is increased cell lysis during the starvation period. Cells which do appear normal, however, often have small polyphosphate bodies. Thus it seems likely that the increasing cell lysis releases phosphate back to the medium, and the surviving cells are able to utilize this phosphate for growth, even at these low phosphate levels.

The two fractions of most interest both during starvation and rapid uptake are the cold TCA extractable and hot TCA extractable phosphates. The cold TCA extract (soluble polyphosphates) decreased by about one-half during the five day starvation period. The hot TCA extract (insoluble polyphosphates) initially is present in lower concentrations than the acid-soluble fraction, but is greater than the acid-soluble fraction at the end of the five day starvation period. The increase in phosphate in the acid-insoluble fraction is also greater after four hours of uptake than in the acid-soluble fraction.

These results are interesting in view of reports of starvation and uptake in other organisms. Aitchison and Butt (1973) reported that phosphate starvation of Chlorella vulgaris resulted in a general reduction of all phosphate fractions, and a 49-90 percent decrease in

the acid-soluble fraction within 8 hours after starvation, and a less rapid but marked decrease in the acid-insoluble fraction. The uptake period was characterized by a rapid increase in all phosphate fractions, with the greatest overcompensation occurring in the acid-soluble fraction. The acid-insoluble fraction only recovered to the levels of cells grown in normal culture. Wiame (1949) found that in yeast, phosphate starvation resulted in a decrease in the insoluble polyphosphate fraction while the soluble polyphosphate remained constant. Phosphate uptake after an induced starvation period led to a large increase in soluble polyphosphate and no appreciable change in the insoluble polyphosphate content. The results of Aitchison and Butt (1973) using Chlorella vulgaris are comparable to those of Wiame (1949).

It has been postulated that the soluble polyphosphates (short chain length) are probably derived from insoluble polyphosphates (long chain length) by the loss of phosphate residues (Krishnan et al, 1957). If this postulate were true, one would expect greater initial increases in soluble polyphosphates during the uptake period, and a later increase in the insoluble polyphosphates due to the addition of phosphate residues to already existing primers, a reversal of the synthesis of short chain polyphosphates.

It is interesting to note that in P. boryanum, the relative levels of acid-soluble and acid-insoluble polyphosphates remain constant throughout the starvation period, even though the levels of acid-insoluble phosphate are lower in normal cells. This would suggest that the acid-soluble fraction is maintained at the expense

of the acid-insoluble fraction. If synthesis occurs by addition of phosphate residues to an already existing primer (acid-soluble fraction), then the amount of acid-insoluble (long chain) phosphates is dependent upon the amount of short chain phosphates and orthophosphate. The greater increase in acid-insoluble polyphosphate at four hours of uptake is probably due to this synthesis from increased short chain primers and available orthophosphate.

Evidence exists which suggests that an enzyme, polyphosphate kinase (ATP: polyphosphate phosphotransferase, E. C. 2.7.4.1., Suzuki et al, 1972) is necessary to polymerize orthophosphate into an osmotically inert form, polyphosphate. This enzyme is synthesized during phosphate starvation (Harold and Harold, 1964; Harold, 1966). Aitchison and Butt (1973) found that in Chlorella vulgaris, polyphosphate synthesis was a consequence of the stimulation of phosphate uptake that was induced by phosphate starvation. These results, along with those of Wiame (1949) lend support to the work of Harold (1966). The synthesis of polyphosphate during rapid uptake is most likely a consequence of an energy dependent uptake of orthophosphate and subsequent polymerization by addition of phosphate residues to pre-existing primers in all organisms reported so far. The polymerization is a result of increased levels of the inducible enzyme, polyphosphate kinase. The results of experiments with P. boryanum are consistent with this hypothesis.

Hydrolysis and Uptake of Condensed Phosphates

The results of hydrolysis studies of sodium tripolyphosphate are presented in Figures 34-36. From these graphs, it is apparent,

especially at the higher phosphate concentrations, that hydrolysis occurs more rapidly and to a greater extent when the algae are present in the medium. At first glance, the graphs are somewhat unclear. Rapid hydrolysis, expressed as orthophosphate present in the medium, increased initially upon transfer, then decreased during the culture period at concentrations of 4.59 and 153.7 mg of phosphate present as condensed phosphate.

These results are consistent with the uptake associated with transfer to fresh medium containing orthophosphate. The algae present in the culture tubes are able to hydrolyze the condensed phosphate, but the resulting orthophosphate is subsequently assimilated by the algae which are already low in total phosphorus levels. Thus there is a reduction in the orthophosphate resulting from hydrolysis. This effect can be seen readily at the lower phosphate concentrations (Figures 34 and 35). The reduction at 1204 mg PO_4 per liter is not evident due to the high phosphate concentration (Figure 36). Hydrolysis of condensed phosphates in the absence of algae is essentially linear.

Figure 37 demonstrates that there is virtually no increase in phosphorus levels in any cell fraction when phosphate-starved algae are inoculated into medium containing condensed phosphates as the phosphorus source. This is probably due to the slow hydrolysis rate of the condensed phosphates. Maximum hydrolysis of the condensed phosphates does not begin to occur until 2 days of culture. Thus, inoculation of phosphorus deficient algae into medium containing condensed phosphates does not result in uptake of phosphate. This

is due to the inability of the algae to assimilate condensed phosphates, and the low levels of orthophosphate present as a consequence of hydrolysis at this early time.

Stewart and Alexander (1971) found that there was an uptake of condensed phosphates present in detergent by the nitrogen-fixing, phosphate starved blue-green alga, Anabaena flos-aquae. However, this uptake occurred over a period of 1 day, and uptake could actually be that of orthophosphate resulting from hydrolysis of the detergent during this time period. The available literature on the utilization of condensed phosphates is scarce. However, it has been demonstrated that hydrolysis of condensed phosphates and assimilation, probably in the form of orthophosphate, occur in several green algae (Davis and Wilcomb, 1967), several blue-green algae (Davis and Wilcomb, 1968), and in the presence of various microorganisms (Clesceri and Lee, 1965a). Hydrolysis is much slower in a sterile environment (Clesceri and Lee, 1965b). Clesceri and Lee (1965b) also found a loss of phosphorus from the medium associated with the hydrolysis of condensed phosphates, probably due to uptake by microorganisms. All these results are consistent with the data presented for P. boryanum.

Electron Microscopy

Cells of P. boryanum grown in different phosphate concentrations exhibit a somewhat different morphology. Cells grown in either 100 or 1000 mg PO_4 per liter culture medium often possess polyphosphate bodies. They also appear to undergo lysis at a much higher frequency. On the other hand, cells grown in 1 mg PO_4 per liter are similar to cells which are grown in phosphate-free medium. They possess large areas

which are of medium electron density, and large intrathylakoidal spaces. Cells grown in 10 mg PO_4 per liter possess inclusions normally found in blue-green algae, and the variations described for the other phosphate concentrations, but to a much smaller degree.

Under conditions of phosphate starvation, the cells develop morphological changes, namely the areas of medium electron density, which later become the areas of polyphosphate body development. These areas may develop at several locations in the cell which have not previously been reported. The first of these is the nucleoplasmic area. This results in the dispersion of DNA toward the periphery of the electron lucent area and subsequent apparent association of DNA with polyphosphate bodies. The electron lucent areas are also observed in the expanded intrathylakoidal spaces, or in polyhedral bodies, with subsequent development of polyphosphate bodies in these areas. The largest polyphosphate bodies are usually located in the nucleoplasmic areas. A few small bodies are usually encountered in phosphate starved cultures. These might be correlated with the fluctuations in phosphorus content of cellular extracts during starvation and the cellular debris, probably resulting from lysis, encountered in sectioned pellets. The polyphosphate bodies could be formed from assimilation of phosphate released during the lysis. The phosphate concentration never approaches that of the external medium, and consequently the polyphosphate bodies formed are much smaller.

The mode of formation of the polyphosphate bodies in the areas of medium electron density and in the polyhedral bodies appears to be

essentially the same as previously reported for the development of polyphosphate bodies in ribosomal areas in Plectonema boryanum (Jensen, 1969). Polyphosphate bodies appear to form in deposition loci, initially an area of medium electron density, and subsequently a "porous body". Electron density increases from the periphery of the body toward the center, until a completely electron dense body is formed (Jensen, 1969). This sequence was also found to be true in this study, regardless of the location of the area of medium electron density.

Thus, in P. boryanum grown under conditions of phosphate starvation and rapid uptake, polyphosphate bodies can develop in 5 different areas of the cell: (1) in ribosomal areas as previously described by Jensen (1969), (2) intrathylakoidally as previously described in Nostoc pruniforme (Jensen, 1968), (3) in association with strands of DNA which is similar to a method previously described by Voelz et al (1966) in Myxococcus xanthus, (4) in areas of medium electron density, and (5) in polyhedral bodies. Voelz et al (1966) described three methods of formation of polyphosphate bodies under different environmental conditions during the "overplus phenomenon" in Myxococcus xanthus. Only the third method described here is similar. Voelz et al (1966) also described dense granules formed around polysaccharide inclusions and dense strands scattered in the cytoplasm.

Another location of polyphosphate body formation has been reported by Dierksheide and Pfister (1973). Under normal culture conditions, they found that in the blue-green alga, Anacystis nidulans,

polyphosphate body formation occurred via the deposition of phosphate within the cyanophycin granule. This site of deposition was never encountered in P. boryanum.

Light microscopy of the cultures demonstrated that there was one apparent polyphosphate body present. Numerous smaller bodies would not be distinct at this limit of resolution. Thus, it is difficult to state with any certainty that one is viewing a single polyphosphate body.

The variation in cell length and the unusual division observed under conditions of phosphate starvation is of interest. Phosphate has been suggested to act as a control factor in cell division in Chlamydomonas reinhardtii (Lien and Knutsen, 1973). A similar type of variation in cell morphology has recently been reported in phosphoglucomutase-deficient mutants of Bacillus licheniformis grown under conditions of phosphate limitation (Forsberg et al, 1973). These mutants of B. licheniformis often developed septa at angles to the normally formed septa, and cell division did not always result in two equal daughter cells.

X-Ray Energy Dispersive Analysis

X-ray energy dispersive analysis of inclusions identified as polyphosphate bodies substantiates earlier physiological data. Polyphosphate bodies, examined under a variety of conditions, contain as the major components phosphorus and calcium. No magnesium is present as a major peak. This does not rule out its presence, but it can be said that it is not a major constituent of polyphosphate bodies in P. boryanum.

At present, there is little available information on x-ray energy dispersive analysis of polyphosphate bodies in other organisms. There are several reports of chemical identification, however, which substantiate the evidence presented here. Rosenberg (1966) found that pyrophosphate granules isolated from Tetrahymena pyriformis contained after ashing 48.5% phosphorus as pyrophosphate, 13.5% calcium, and 8.1% magnesium by weight. Munk and Rosenberg (1969) investigating this same organism, found that equimolar concentrations of calcium and magnesium ions were essential for the deposition of pyrophosphate granules. Isolated granules were found to be deposits of calcium magnesium pyrophosphate. Friedberg and Avigad (1968) reported that polyphosphate bodies isolated from Micrococcus lysodeikticus contained 18.39 ug Mg per mg dry weight and 7.54 ug Ca per mg dry weight. Cytoplasmic granules of Tetrahymena pyriformis analyzed in situ by electron probe analysis (Coleman et al, 1972) were found to contain potassium, calcium, magnesium, phosphorus, and a lipid-like material. The ratios of calcium to phosphorus and magnesium to phosphorus tended to divide the granules into two distinct classes. These reports are in some agreement with the data presented here.

It can be seen that much work remains to be done in this area. At present, no exhaustive study has been conducted previously on one organism, making a comparison of these studies with those reported in the literature difficult to interpret. The literature on phosphate metabolism is composed of a variety of studies utilizing diverse organisms as well as a variety of environmental and physiological

parameters. Thus, selection of one organism to study seems essential.

The relationship between the laboratory investigations reported here and the field observations of other workers is also of interest. It has been suggested that phosphorus is one of the key elements implicated in the limiting nutrient controversy. The studies with Plectonema boryanum indicate that cultures quite easily become phosphorus limited. When put into a variety of situations where phosphorus is available as orthophosphate, the algae can easily assimilate large quantities of phosphate and store it as both soluble (short chain) and insoluble (long chain) polyphosphates in the cellular inclusion known as a polyphosphate body. This then probably serves as a phosphorus reserve. The cells are still able to grow when phosphorus appears to be limiting in the medium.

Studies with uptake of condensed phosphates do not rule out detergents as sources of phosphate which are able to support the growth of P. boryanum. They merely indicate that condensed phosphates cannot be utilized directly, but must first be hydrolyzed to orthophosphate. The orthophosphate can then be assimilated and used for synthesis. It is most probable that algae, in their normal environment, are phosphorus deficient. Possession of a mechanism whereby phosphate entering the environment could be assimilated rapidly and stored in an osmotically inert form would enable organisms to survive in periods of low available phosphorus. Phosphorus could enter the environment, for example, through sewage, or land runoff after a rainfall. It is suggested that measurement of phosphate concentrations in natural bodies of water should not be the criterion for

establishing phosphate levels which are conducive to accelerating eutrophication. A more meaningful estimation would be the total phosphorus present in the cell, since phosphorus stored in the cell as polyphosphate is capable of sustaining growth, even when there is no apparent available phosphate in the environment.

SUMMARY

Physiological and cytological aspects of phosphate utilization by the blue-green alga *Blectonema boryanum* were studied. It was found that the external phosphate concentration influenced the distribution of phosphorus-containing compounds as well as cell architecture.

Culturing the alga in concentrations of 100, and 1000 mg PO_4 per liter resulted in increases in the levels of acid-insoluble and acid soluble polyphosphates. The values reported for 100 and 1000 mg PO_4 /liter were the same, indicating that the algae were able to assimilate and utilize only fixed amounts of phosphates. This value was calculated to be 6.5 ug P per 10^6 cells. Increased external phosphate concentration led to increases in size and in the frequency of encountering polyphosphate bodies in the cells. However, cell death and lysis also increased at these higher concentrations.

When the algae were grown in a concentration of 1 mg PO_4 /liter, the phosphate present in all cell fractions decreased. Cytologically, the cell was also altered. Large areas of medium electron density developed, and the area of intrathylakoidal spaces increased.

The changes that were encountered in the cells grown in 1 mg PO_4 /liter were similar but reduced, when compared to algae grown in phosphate-free medium for 5 days. It was determined that cells grown in the absence of phosphate for 5 days had total cell phosphorus levels of 0.76 ug P per 10^6 cells. Cells in culture for two months or longer were found to have total cell phosphorus levels of 0.73 ug P/ 10^6

cells. This was determined to be the minimum cell phosphorus level, limiting growth. Transfer of cells from either of the two culture conditions previously described to a medium containing phosphate led to an "overplus" phenomenon.

The phosphate overcompensation reaction was characterized by increases in all cell phosphorus fractions. The most dramatic increase was that of both soluble and insoluble polyphosphates. These fractions often increased by more than an order of magnitude. The greatest phosphate uptake occurs within one hour of transfer of phosphate starved cells into a medium containing a known amount of phosphate, and is essentially complete at 4 hours. Phosphate uptake is light dependent and is inhibited by 2,4-dinitrophenol, mercuric chloride, and sodium fluoride. It is suggested that the energy required for the uptake process is the result of the metabolism of photosynthetic products. The total cell phosphorus levels for uptake never increased beyond $18.9 \text{ ug P}/10^6 \text{ cells}$.

The phosphate overplus phenomenon was also quite distinct cytologically. Large polyphosphate bodies developed in the areas of medium electron density. These areas developed during phosphate-free or phosphate-limited growth. The areas of medium electron density, and consequently, polyphosphate bodies developed in 5 different locations in the cell: 1. In ribosomal areas, 2. intrathylakoidally, 3. in nucleoplasmic areas, 4. in polyhedral bodies, and 5. in apparent association with strands of DNA. All developmental stages of polyphosphate bodies can be found during this uptake process.

It was also determined that the alga could not utilize sodium tripolyphosphate, a condensed linear phosphate, for the uptake phenomenon. It appeared that hydrolysis of the condensed phosphate was essential for subsequent utilization by the alga.

X-ray energy dispersive analysis of polyphosphate bodies revealed that there were two major components of a polyphosphate body--phosphorus and calcium. Magnesium or other elements were not detectable in amounts that are greater than background emission.

APPENDICES

Appendix A

Composition of Culture Medium - Modified Fitzgerald (Fitzgerald et al, 1952)

Concentration is expressed in mg/liter

NaNO ₃	-	124
K ₂ HPO ₄ · 3H ₂ O	-	13
MgSO ₄ · 7H ₂ O	-	25
CaCl ₂ · 2H ₂ O	-	36
Na ₂ CO ₃	-	20
Na ₂ SiO ₃ · 9H ₂ O	-	58
Ferric Citrate	-	3
Citric Acid	-	3
Gaffron's minor element solution - 0.04ml		

Gaffron's solution. (in g/liter)

H ₃ BO ₃	-	3.10
MnSO ₄ · 4H ₂ O	-	2.23
ZnSO ₄ · 7H ₂ O	-	0.287
(NH ₄) ₆ Mo ₇ O ₂₄ · 4H ₂ O	-	0.088
CuSO ₄ · 5H ₂ O	-	0.125
Co(NO ₃) ₂ · 6H ₂ O	-	0.146
Al ₂ (SO ₄) ₃ · K ₂ SO ₄ · 24H ₂ O	-	0.474
NiSO ₄ (NH ₄) ₂ SO ₄ · 6H ₂ O	-	0.198

$\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$	- 0.154
$\text{Cr}(\text{NO}_3)_3 \cdot 7\text{H}_2\text{O}$	- 0.037
$\text{V}_2\text{O}_5(\text{SO}_4)_3 \cdot 16\text{H}_2\text{O}$	- 0.035
$\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$	- 0.033
KBr	- 0.119
KI	- 0.083

Appendix B

Reagents for Murphy-Riley Orthophosphate Determination (Murphy and Riley, 1962)

1. Sulfuric acid (5N). Dilute 70 ml of concentrated reagent grade sulfuric acid to 500 ml.
2. Ammonium molybdate. Dissolve 20 grams of reagent grade ammonium molybdate in water and dilute to 500 ml. Store the solution in a Pyrex glass bottle.
3. Ascorbic acid (0.1M). Dissolve 1.32 grams of ascorbic acid in 75 ml of distilled water. This solution is made on the day it is required.
4. Potassium antimonyl tartrate (1mg Sb/ml). Dissolve 0.2743 grams of potassium antimonyl tartrate in distilled water and dilute to 100 ml.
5. Mixed reagent. Mix thoroughly 125 ml of 5N sulfuric acid and 37.5 ml of ammonium molybdate. Add 75 ml of ascorbic acid solution and 12.5 ml of potassium antimonyl tartrate solution. This reagent keeps for 24 hours.

Appendix C

Microscopy Procedures

Schedule 1 Preparation of solutions

1. Stock Michaelis Buffer.

a. Weight: 1.94 grams sodium acetate ($\text{NaC}_2\text{H}_3\text{O}_2 \cdot 3\text{H}_2\text{O}$)

2.94 grams sodium barbital

3.40 grams sodium chloride

b. Add water to 100 ml

c. Store in refrigerator

2. Cacodylate buffer.

A stock solution of 0.2M cacodylate buffer was made by dissolving 42.8 grams of sodium cacodylate $\text{Na}(\text{CH}_3)_2\text{AsO}_2 \cdot 3\text{H}_2\text{O}$ in 1 liter of distilled water. The pH was adjusted to pH 6.2 before the solution was brought to final volume.

3. Osmium Tetroxide

A stock solution of 2% OsO_4 was made by dissolving 1 gram of osmium tetroxide in 50 ml of glass distilled water. The solution was stored in a Pyrex glass stoppered bottle and stored in the refrigerator.

4. 3% Glutaraldehyde

Fresh solutions of 3% glutaraldehyde in 0.1M cacodylate buffer were made by mixing 3 ml of 50% biological grade glutaraldehyde, 22 ml glass distilled water, and 25 ml of 0.2M cacodylate buffer.

5. 1% Bacto-tryptone

1 gram of Difco bacto-tryptone and 0.5 grams NaCl were

dissolved in 100 ml glass distilled water. The solution was placed in small closed vials and stored frozen. For use, the contents of the vials were thawed and mixed thoroughly before use.

6. Epon 812 (Luft, 1961)

Mixture A.

62 ml Epon 812

100 ml DDSA (Dodecanyl Succinic Anhydride).

Mixture B

100 ml Epon 812

89 ml NMA (Nadic Methyl Anhydride).

1 part mixture A, and 2 parts mixture B were mixed in a disposable plastic beaker, and 0.2 ml of DMP-30 was added per 10 ml Epon mixture. It is important that all components are thoroughly mixed so that no "streaks" are visible. Stock solutions of Mixtures A and B were refrigerated, and brought to room temperature before opening to avoid condensation.

Schedule 2 - Fixing and Embedding Procedures

A. Modified Kellenberger (Modified Osmium) (Penkratz and Bowen, 1963).

(1) 5 ml Stock Michaelis Buffer

7 ml 0.1 N HCl

13 ml distilled water

0.25 ml 1.0M CaCl₂

adjust pH to 6.1 or 6.2

(2) Dilute (1.) 1:1 with 2% OsO₄

Add 0.1 ml 1% bacto-tryptone per ml fixative (2).

(3) Fix for 3 hours at room temperature.

B. Dehydration

After fixation, the algae were dehydrated and embedded in Epon 812 as follows:

50% EtOH - 5 minutes

70% EtOH - 5 minutes

85% EtOH - 5 minutes

95% EtOH - 5 minutes

100% EtOH - 5 minutes

100% EtOH - 5 minutes

100% EtOH - 5 minutes

Propylene Oxide - 5 minutes

Propylene Oxide - 5 minutes

Propylene Oxide - 5 minutes

Propylene Oxide: Epon (3:1) - 1 hour

Propylene Oxide: Epon (1:1) - 1 hour

Pure Epon - overnight

Samples were pelleted into Beem capsules and the epoxy polymerized as follows:

35°C (Incandescent lamp) - 1 day.

45°C - 1 day.

65°C - 2 days.

C. Preparation of post stains

1. Uranyl acetate - methanol stain (Stempak and Ward, 1964).

a. Preparation 15 grams of hydrated uranyl acetate $(UO_2(CH_3COO)_2 \cdot 2H_2O)$ were dissolved in 50 ml of acetone-free absolute methanol with a magnetic stirrer. The solution was then stored in a Pyrex glass-stoppered bottle at 4°C.

b. Staining

Grids were immersed in wells of a porcelain staining dish, section side up, for 10 minutes at room temperature. The wells were covered during this period to prevent evaporation of the methanol. The grids were then rinsed with absolute methanol, absolute ethanol, and water.

2. Lead - citrate stain (Reynolds, 1963)

a. Preparation 1.33 grams of lead nitrate $(Pb(NO_3)_2)$ and 1.76 grams of sodium citrate $(Na_3(C_6H_5O_7) \cdot 2H_2O)$ were added to 30 ml of fresh glass distilled water in a 50 ml volumetric flask. The flask was shaken vigorously for one minute, and intermittently for $\frac{1}{2}$ hour. 80 ml of 1N NaOH was then added and mixed. The solution was then brought to volume with distilled water. This solution was stored in a glass-stoppered volumetric flask at room temperature.

b. Staining

Grids were floated on drops of the stain on parafilm in petri dishes containing water and sodium hydroxide pellets for 10 minutes. The grids were then washed

with a fresh solution of 0.02 N NaOH followed by distilled water.

**D. Preparation of Fluka Durcupan Water-Soluble Embedding Medium
(Polysciences, Fluka Durcupan embedding kit)**

a. Components

"A" - a water soluble aliphatic polyepoxide

"B" - 964 hardener, an anhydride of a diazide with aliphatic side chain.

"C" - 960 hardener, a phenol derivative with amino groups.

"D" - plasticizer (dibutyl phthalate)

b. Dehydration Bath

1. 50% "A" + 50% water - 30 min.
2. 70% "A" + 30% water - 45 min.
3. 90% "A" + 10% water - 45 min.
4. 100% "A" - 90 min.
5. 100% "A" - 90 min.

c. Polymerization Mixture

"A" - 5 ml

"B" - 11.7 ml

"C" - 1.1 ml

"D" - 0.3 ml

- d. Embedding** - Leave tissue overnight at 4-5°C. Pour fresh polymerization mixture into Beem capsules and add samples. Polymerize 37-45°C for 3-4 days.

LITERATURE CITED

- Aitchison, P. A., and V. S. Butt. 1973. The relation between the synthesis of inorganic polyphosphate and phosphate uptake by Chlorella vulgaris. Journal of Experimental Botany 24:497-510.
- American Public Health Association. 1965. Standard methods for the examination of water and wastewater. 12th ed. American Public Health Association. 626pp.
- Atkins, W. R. G. 1923. The phosphate content of fresh and salt waters in its relationship to the growth of the algal plankton. Journal of the Marine Biological Association, United Kingdom 13:119-150.
- Atkins, W. R. G. 1925. Seasonal changes in the phosphate content of sea water in relation to the growth of the algal plankton during 1923 and 1924. Journal of the Marine Biological Association, United Kingdom 13:700-720.
- Baker, A. I., and R. R. Schmidt. 1964. Further studies on the intracellular distribution of phosphorus during synchronous growth of Chlorella pyrenoidosa. Biochimica et Biophysica Acta 82:336-342.
- Batterton J. C., and C. Van Baalen. 1968. Phosphorus deficiency and phosphate uptake in the blue-green alga, Anacystis nidulans. Canadian Journal of Microbiology 14:341-348.
- Belsky, M. M., S. Goldstein, and Maria Menna. 1970. Factors affecting phosphate uptake in the marine fungus Dermocystidium sp. Journal of General Microbiology 62:399-402.

- Blum, J. J. 1966. Phosphate uptake by phosphate-starved Euglena.
Journal of General Physiology 49:1125-1136.
- Bone, Derek H. 1971. Relationship between phosphates and alkaline
phosphatase of Anabaena flos-aquae in continuous culture.
Archives fur Mikrobiologie 80:147-153.
- Borst-Pauwels, G. W. F. H., and S. Jager. 1969. Inhibition of phosphate
and arsenate uptake in yeast by monoiodoacetate, fluoride,
2,4-dinitrophenol, and acetate. Biochimica et Biophysica Acta
172:399-406.
- Bringmann, G. 1950. Vergleichende licht-und elektronen-mikroskopische
Untersuchungen an Oszillatorien. Planta 38:541.
- Bush, A. F., and S. F. Mulford. 1954. Studies of Waste Water Reclama-
tion and Utilization. California State Water Pollution
Control Board, Sacramento, Pub. No. 9, 82pp.
- Butt, W. D., and H. Lees. 1960. The biochemistry of nitrifying
organisms. Part 7. The phosphate compounds of Nitrobacter
and the uptake of orthophosphate by the organism. Canadian
Journal of Biochem. Physiol. 38:1295-1300.
- Cashe, Michael, and Ernst Freese. 1964. Excretion of alkaline
phosphatase by Bacillus subtilis. Biochemical and Biophysical
Research Communications 16:541-544.
- Chayen, R., Sonya Chayen, and E. R. Roberts. 1955. Observations on
nucleic acid and polyphosphate in Torulopsis utilis.
Biochimica et Biophysica Acta 16:117-126.

- Clesceri, Nicholas L., and G. Fred Lee. 1965a. Hydrolysis of Condensed Phosphates I. Non-sterile environment. International Journal of Air and Water Pollution 9:723-742.
- Clesceri, Nicholas L., and G. Fred Lee. 1965b. Hydrolysis of Condensed Phosphates II. Sterile environment. International Journal of Air and Water Pollution 9:743-751.
- Cole, J. A., and D. E. Hughes. 1965. The metabolism of polyphosphates in Chlorobium thiosulfatophilum. Journal of General Microbiology 38:65-72.
- Coleman, J. R., Jytle R. Nilsson, R. R. Warner, and P. Batt. 1972. Qualitative and Quantitative Electron Probe analysis of cytoplasmic granules in Tetrahymena pyriformis. Experimental Cell Research 74:207-219.
- Cooke, G. W., and R. J. B. Williams. 1973. Significance of man-made sources of phosphorus: fertilizers and farming. The phosphorus involved in agricultural systems and possibilities of its movement into natural water. Water Research 7:19-34.
- Correll, David L. 1965. Ribonucleic acid-polyphosphate from algae. III. Hydrolysis studies. Plant and Cell Physiology 6:661-669.
- Correll, David L., and N. E. Tolbert. 1962. Ribonucleic acid-polyphosphate from algae. I. Isolation and physiology. Plant Physiology 37:627-636.
- Correll, David L., and N. E. Tolbert. 1964. Ribonucleic acid-polyphosphate from algae. II. Physical and chemical properties of the isolated complexes. Plant and Cell Physiology 5:171-191.

- Cowen, W., and G. Fred Lee. 1971. Leaves as a source of phosphorus. Report Water Chemistry Program University of Wisconsin. Madison, Wisconsin.
- Davis, Ernst M., and Maxwell J. Wilcomb. 1967. Enzymatic degradation and assimilation of condensed phosphates by green algae. Water Research 1:335-350.
- Davis, Ernst M., and Maxwell J. Wilcomb. 1968. Condensed phosphate degradation and assimilation by selected blue-green algae. Water Research 2:311-324.
- Devey, D. G., and N. Harkness. 1973. The significance of man-made sources of phosphorus: detergents and sewage. Water Research 7:35-54.
- Dierksheide, William C., and Robert M. Pfister, 1973. Associated organelles in the blue-green alga, Anacystis nidulans. Canadian Journal of Microbiology 19:149-151.
- Drews, G. 1960. Elektronenmikroskopische untersuchungen an Mycobacterium phlei. Archives fur Mikrobiologie 35:53-62.
- Drews, G., and W. Niklowitz. 1957. Beitrage zur Cytologie der Blaualgen. III. Untersuchungen uber die granularen Einschlusse der Hormogonales. Archives fur Mikrobiologie 25:333-351.
- Duguid, J. P. 1948. The influence of cultural conditions on the morphology of Bacterium aerogenes with reference to nuclear bodies and capsule size. Journal of Pathology and Bacteriology 60:265-274.

- Duguid, J. P., I. W. Smith, and J. F. Wilkinson. 1954. Volutin production in Bacterium aerogenes due to development of an acid reaction. Journal of Pathology and Bacteriology 62:289-300.
- Ebel, J. P., A. Stair, G. Dirheimer, S. Muller-Felter, and M. Yacoub. 1962. Relations de structure entre acides ribonucleiques et polyphosphates. Côllôq. Intern. Centre Natl. Rech. Sci. (Paris) 106:545-573.
- Ebel, J. P., and J. Colas, and S. Muller. 1958a. Recherches cytochimiques sur les polyphosphates. II. Mise au point de methodes de detection cytochimiques specifiques des polyphosphates. Experimental Cell Research 15:28-36.
- Ebel, J. P., J. Colas, et S. Muller. 1958b. Recherches cytochimiques sur les polyphosphates imorganiques contenus dans les organismes vivant. III. Presence de polyphosphates chez divers organismes inferieurs. Experimental Cell Research 15:36-42.
- Edmunson, W. T. 1961. Changes in Lake Washington following an increase in the nutrient income. Verh. Int. Ver. Limnol. 14:167-175.
- Edmunson, W. T. 1969. Cultural eutrophication with special reference to Lake Washington. Mitt. Int. Verein. Limnol. 17:19-32.
- Edmunson, W. T. 1970. Phosphorus, nitrogen, and algae in Lake Washington after diversion of sewage. Science 169:690-691.

- Edmunson, W. T. 1972. Nutrients and Phytoplankton in Lake Washington. In: Nutrients and Eutrophication: The Limiting Nutrient Controversy. G. E. Likens, ed. American Society of Limnology and Oceanography, Special Symposia, Volume 1. pp 172-196.
- Engelbrecht, Richard S., and James J. Morgan. 1959. Studies on the occurrence and degradation of condensed phosphate in surface waters. Sewage and Industrial Wastes 31:458-478.
- Ennor, A. H., and J. F. Morrison. 1958. Biochemistry of the phosphagens and related guanidines. Physiological Reviews 38:631-674.
- Eppley, R. W. 1962. Hydrolysis of polyphosphates by Porphyra and other seaweeds. Physiologia Plantarum 15:246-251.
- Ferguson, A. 1968. A nonmyopic approach to the problems of excess algae growths. Environmental Science and Technology 2:188.
- Fippen, E. O. 1945. Plant nutrient losses in silt and water in the Tennessee river system. Soil Science 60:223.
- Fisher, Knute A. 1971. Polyphosphate in a chlorococcalean alga. Phycologia 10:177-182.
- Fitzgerald, George P. 1970. Evaluations of the availability of sources of nitrogen and phosphorus for algae. Journal of Phycology 6:239-247.
- Fitzgerald, G. P., and T. C. Nelson. 1966. Extractive and enzymatic analysis for limiting or surplus phosphorus in algae. Journal of Phycology 2:32-37.

- Fitzgerald, G. P., and G. A. Rohlich. 1964. Biological removal of nutrients from sewage; laboratory experiments. Verh. int. Verein. theor. Angew. Limnol. 15:597-608.
- Fitzgerald, G. P., G. C. Gerloff, and F. Skoog. 1952. Studies on chemicals with selective toxicity to blue-green algae. Sewage and Industrial Wastes 24:888-896.
- Fogg, G. E. 1973. Phosphorus in primary aquatic plants. Water Research 7:77-92.
- Forsberg, E. W., P. B. Wyrick, J. B. Ward, and H. J. Rogers. 1973. Effect of phosphate limitation on the morphology and wall composition of Bacillus lichiniformis and its phosphoglucomutase-deficient mutants. Journal of Bacteriology 113:969-984.
- Friedberg, Ilan, and Gad Avigad. 1968. Structures containing polyphosphate in Micrococcus lysodeikticus. Journal of Bacteriology 96: 544-553.
- Fuhs, G. W. 1958. Uber die natur der granula im Cytoplasma von Oscillatoria amoena (Kutz) Gom. Osterreich Bot. 2. 104:531-551.
- Fuhs, G. Wolfgang. 1969. The Nuclear Structures of Protocaryotic Organisms (Bacteria and Cyanophyceae). Protoplasmatologia Vol. 4 186 pp. Springer - Verlag. N. Y.
- Fuhs, G. Wolfgang, Suzanne D. Demmerle, Edmondo Canelli, and Min Chen. 1972. Characterization of Phosphorus - Limited Plankton Algae (With reflections on the limiting-nutrient concept) In: Nutrients and Eutrophication: The Limiting Nutrient Contro-

- versy. G. E. Likens, ed. American Society of Limnology and Oceanography, Special Symposia, Volume 1. pp 113-133.
- Galloway, R. A., and R. W. Krauss (Editors) 1963. Utilization of phosphorus sources by Chlorella. In: Studies of Microalgae and Photosynthetic Bacteria, pp. 569-575. Japanese Soc. of Plant Physiol. University of Tokyo, Tokyo, Japan.
- Gibson, C. E. 1971. Nutrient Limitation. Journal of the Water Pollution Control Federation 43:2436-2440.
- Golterman, H. L. 1973. Natural phosphate sources in relation to phosphate budgets: A contribution to the understanding of eutrophication. Water Research 7:3-18.
- Grula, Edward A., R. H. Weaver, and O. F. Edwards. 1954. Studies on a strain of Caulobacter from water. II. Nutrition, with implications for cytology. Journal of Bacteriology 68:201-206.
- Harold, F. M. 1960. Accumulation of inorganic polyphosphate in mutants of Neurospora crassa. Biochimica et Biophysica Acta 45:172-188.
- Harold, F. M. 1962. Depletion and replenishment of the inorganic polyphosphate pool in Neurospora crassa. Journal of Bacteriology 83:1047-1057.
- Harold, F. M. 1963. Accumulation of inorganic polyphosphate in Aerobacter aerogenes. I. Relationship to growth and nucleic acid synthesis. Journal of Bacteriology 86:216-221.
- Harold, F. M. 1964. Enzymic and genetic control of polyphosphate accumulation in Aerobacter aerogenes. Journal of General Microbiology 35:81-90.

- Harold, F. M. 1965. Regulatory mechanisms in the metabolism of inorganic polyphosphate in Aerobacter aerogenes. Colloq. Intern. Centre Natl. Rech. Sci. (Paris) 124, pp. 307-315.
- Harold, Franklin M. 1966. Inorganic polyphosphates in biology: structure, metabolism, and function. Bacteriological Reviews 30:772-794.
- Harold, F. M., and R. L. Harold, 1965. Degradation of inorganic polyphosphate in mutants of Aerobacter aerogenes. Journal of Bacteriology 89:1262-1270.
- Harold, R. L., and F. M. Harold. 1963. Mutants of Aerobacter aerogenes blocked in the accumulation of inorganic polyphosphate. Journal of General Microbiology 31:241-246.
- Harold, F. M., and S. Sylvan. 1963. Accumulation of inorganic polyphosphate in Aerobacter aerogenes. II. Environmental control and the role of sulfur compounds. Journal of Bacteriology 86:222-231.
- Harris, E., and G. A. Riley. 1956. Oceanography of Long Island Sound. VIII. Chemical composition of the plankton. Bull. Bingham Oceanogr. Collect. Yale University 15:315-323.
- Harwood, J. E., R. A. Van Steenderen and A. L. Kuhl. 1969. A comparison of some methods for total phosphate analysis. Water Research 3:425-432.
- Hasler, A. D. 1947. Eutrophication of lakes by domestic drainage. Ecology 28:383-395.
- Hosokawa, Iwao, and Fumio Ohshima. 1973. An improved method of phosphorus analysis in sea water. Water Research 7:283-290.

- Hudson, E. J., and H. W. Marson. 1970. Eutrophication: with particular reference to the role of phosphates. Chemistry and Industry 4:1449-1458.
- Huennekens, F. M., and H. R. Whiteley. 1960. Phosphoric acid anhydrides and other energy rich compounds. In: Comparative Biochemistry. Vol. 1. Ed. M. Florkin and H. S. Mason pp. 107-180. Academic Press, N. Y.
- Hughes, D. E., and A. Muhammed. 1962. The metabolism of polyphosphate in bacteria. Colloq. Int. Centre Nat. Rech. Sci. 106:591-602.
- Hutchinson, G. E. 1957. A treatise on Limnology. Vol. I Geography, Physics, and Chemistry. J. Wiley & Sons, New York. 1015 pp.
- Hutchinson, G. Evelyn. 1973. Eutrophication. American Scientist 61:269-279.
- Indge, K. J. 1968. Polyphosphates of the yeast cell vacuole. Journal of General Microbiology 51:447-455.
- Jeener, R., and J. Brachet. 1944. Recherches sur l'acide ribonucleique des levures. (Microdosage, relations avec la croissance, conditions de sa synthese.) Enzymologia 11:222-234.
- Jenkins, David, Warren J. Kaufman, P. H. McGauhey, Alex J. Horne, and James Gasser, 1973. Environmental impact of detergent builders in California waters. Water Research 7:265-281.
- Jensen, Thomas E. 1968. Electron microscopy of polyphosphate bodies in a blue-green alga, Nostoc pruniforme. Archives fur Mikrobiologie 62:144-152.
- Jensen, Thomas E. 1969. Fine structure of developing polyphosphate bodies in a blue-green alga, Plectonema boryanum. Archives

- fur Mikrobiologie 67:328-338.
- Juni, E., M. O. Kamen, S. Spiegelman, and J. M. Wiame. 1947.
Physiological Heterogeneity of Metaphosphate in Yeast.
Nature 160:717-718.
- Jungnickel, F. 1970. Untersuchungen zum einfluß des 2,4-Dinitrophenols
auf Phosphat-und Kaliumstoffwechsel phosphatverarmter Zellen
von Candide utilis. Zentralblatt fur Bakteriologie,
Parasitenkunde, Infektionskrankheiten und Hygiene. 125:243-249.
- Kaltwasser, H. 1962. Die rolle der polyphosphate im Phosphatstoffioecasel
eines Knallgasbakteriums (Hydrogenomonas Stamm 20). Archives
fur Mikrobiologie 41:282-306.
- Kanai, Ryuzi, Shigeji Aoki, and Shigetoh Miyachi. 1965. Quantitative
separation in inorganic polyphosphates in Chlorella cells.
Plant and Cell Physiology 6:467-473.
- Kanai, R., S. Miyachi, and S. Miyachi 1963. Light-induced formation
and mobilization of polyphosphate "C" in Chlorella cells.
Plant Cell Physiol. (Tokyo) special issue. Studies on
microalgae and photosynthetic bacteria. pp. 613-618.
- Kanai, R., and W. Simonis. 1968. Einbau von ³²P in Verschiedene
Phosphatfraktionen, besonders Polyphosphate, bei einzelliger
Grunalgen (Ankistrodesmus braunii) im Licht und im Dunkeln.
Archives fur Mikrobiologie 62:56-71.
- Katchman, Bernard G., and William O. Fetty. 1955. Phosphorus
metabolism in growing cultures of Saccharomyces cerevisiae.
Journal of Bacteriology 69:607-615.

- Katchman, Bernard J., and John R. Van Wazer. 1954. The "soluble" and "insoluble" polyphosphates of yeast. Biochimica et Biophysica Acta 14:445-446.
- Ketchum, Bostwick H. 1939. The development and restoration of deficiencies in the phosphorus and nitrogen composition of unicellular plants. Journal of Cellular and Comparative Physiology 13:373-381.
- Ketchum, Bostwick H., 1966. The absorption of phosphate and nitrate by illuminated cultures of Nitzschia closterium. American Journal of Botany 26:399-407.
- Keup, Lowell E. 1968. Phosphorus in Flowing Waters. Water Research 2:373-386.
- Kholy, A. 1956. On the assimilation of phosphorus in Chlorella pyrenoidosa. Physiologie Plantarum 9:137-143.
- Knaysi, G., and R. F. Baker. 1947. Demonstration, with the electron microscope, of a nucleus in Bacillus mycoides grown in a nitrogen-free medium. Journal of Bacteriology 53:539.
- Knaysi, G., J. Hillier, and C. Fabricant. 1951. The cytology of an avian strain of Mycobacterium tuberculosis studied with the electron and light microscope. Journal of Bacteriology 60:423.
- Knaysi, Georges and Stuart Mudd. 1943. The internal structure of certain bacteria as revealed by the electron microscope. A contribution to the study of the bacterial nucleus. Journal of Bacteriology 45:349-359.

- Konig, H., and A. Winkler. 1948. Uber Einschlusse in Bakterien und ihre Veranderung in Elektronenmikroskop. Naturwissenschaften 35:136-144.
- Kornberg, A., S. R. Kornberg, and E. S. Simms. 1956. Metaphosphate synthesis by an enzyme from Escherichia coli. Biochimica et Biophysica Acta 26:215-227.
- Kornberg, S. R. 1956. Tripolyphosphate and trimetaphosphate in yeast extracts. Journal of Biological Chemistry 218:23-31.
- Kornberg, S. R. 1957. Adenosine triphosphate synthesis from polyphosphate by an enzyme from Escherichia coli. Biochimica et Biophysica Acta 26:294-300.
- Krishnan, P. S., S. P. Damle, and Violet Bajaj. 1957. Studies on the role of "metaphosphate" in molds. II. The formation of "soluble" and "insoluble" metaphosphates in Aspergillus niger. Archives of Biochemistry and Biophysics 67:35-52.
- Kuentzel, L. E. 1969. Bacteria, Carbon Dioxide, and Algal Blooms. Journal of the Water Pollution Control Federation 41:1737-1747.
- Kuenzler, E. J., and B. H. Ketchum. 1962. Rate of phosphorus uptake by Phaeodactylum tricornutum. Biological Bulletin 123:134-145.
- Kuhl, A. 1968. Phosphate metabolism of green algae. In: Algae, Man, and the Environment ed. D. F. Jackson, pp. 37-52. Syracuse University Press.
- Kulaev, I. S., and A. N. Belozerskii. 1958. Electrophoretic studies on polyphosphate-ribonucleic acid complexes from Aspergillus niger. Proc. Acad. Sci. USSR (English Transl.) 120:128-131.

- Kylin, Anders. 1966. The influence of photosynthetic factors and metabolic inhibitors on the uptake of phosphate in P-deficient Scenedesmus. Physiologia Plantarum 19:644-649.
- Langen, P. 1965. Vorkommen und Bedeutung von Polyphosphaten in Organismen. Biol. Rundschau. 2:145-152.
- Langen, P., and E. Liss. 1958. Über Bildung und Umsatz der Polyphosphate der Hefe. Biochem. Z. 330:455-466.
- Lean, D. R. S. 1973. Phosphorus dynamics in lake water. Science 179:678-679.
- Lembke, A., and H. Ruska. 1940. Vergleichende mikroskopische und Übermikroskopische Beobachtungen an den Irregesen der Tuberkulose. Klin. Wochenschi 19:217-220.
- LePage, G. A. 1949. Methods for the analysis of phosphorylated intermediates, p. 190. In: Manometric techniques and related methods for the study of tissue metabolism. Edited by W. W. Umbreit, R. H. Burris, and J. F. Stauffer. Burgess Publishing Co. Minneapolis, Minnesota.
- Levin, Gilbert V., and Joseph Shapiro 1965. Metabolic uptake of phosphorus by wastewater organisms. Journ. Water Poll. Cont. Fed. 37:800-821.
- Lewin, Valentine H. 1973. Phosphates in sewage and sewage treatment. Water Research. 7:55-68.
- Lien, T., and G. Knutsen. 1973. Phosphate as a control factor in cell division of Chlamydomonas reinhardtii, studied in synchronous culture. Experimental Cell Research 78:70-88.

- Likens, G. E. 1972. Eutrophication and aquatic ecosystems. In: Nutrients and Eutrophication: The Limiting Nutrient Controversy. G. E. Likens, Ed. American Society of Limnology and Oceanography, Special Symposia, Volume 1. pp. 3-13. Lawrence, Kansas.
- Likens, G. E. (ed.) 1972. Nutrients and Eutrophication: The Limiting Nutrient Controversy. American Society of Limnology and Oceanography. Special Symposia, Volume 1. Lawrence, Kansas.
- Lindgren, Carl. C. 1948. The origin of volutin on the chromosomes, its transfer to the nucleolus, and suggestions concerning the significance of this phenomenon. Proceedings of the National Academy of Science 34:187-193.
- Liss, E., and P. Langen. 1962. Versuche zur polyphosphat-überkompensation in hefezellen nach phosphatverarmung. Archives für Mikrobiologie 41:383-392.
- Luft, J. H. 1961. Improvements in epoxy resin embedding methods. Journal of Biophysical and Biochemical Cytology. 9:409-414.
- Macfarlane, Marjorie Giffen. 1936. Phosphorylation in living yeast. Biochemical Journal 30:1369-1379.
- Mackereth, F. J. 1953. Phosphorus utilization by Asterionella formosa Hass. Journal of Experimental Botany 4:296-313.
- Marson, Harold. 1971. The removal of phosphates from sewage. Effluent Water Treatment Journal 1:309-315.
- Menzil, D. W., and N. Corwin. 1965. The measurement of total phosphorus in seawater based on the liberation of organically bound fractions by persulfate oxidation. Limnology and Oceanography 10:280-282.

- Meyer, A. 1904. Orientierende Untersuchungen über Verbreitung, Morphologie and Chemie des Volutins. Bot. Ztg. 62:113.
- Mitchell, Dee. 1971. Eutrophication of Lake Water Microcosms: Phosphate versus nonphosphate Detergents. Science 174:827-829.
- Miyachi, S., R. Kanai, S. Mihara, S. Miyachi, and S. Aoki. 1964. Metabolic roles of inorganic polyphosphates in Chlorella cells. Biochimica et Biophysica Acta 93:625-634.
- Miyachi, S., and H. Tamiya. 1961. Distribution and turnover of phosphate compounds in growing Chlorella cells. Plant and Cell Physiology 2:405-414.
- Morton, H. E., and T. F. Anderson. 1941. Electron microscopic studies of biological reactions. I. Reduction of potassium tellurite by Conyebacterium diphtheriae. Proceedings of the Society for Experimental Biology and Medicine 46:272-276.
- Mudd, S. 1953. The mitochondria of bacteria. Journal of Histochemistry and Cytochemistry 1:248-253.
- Mudd, S., A. F. Brodie, L. C. Winterscheid, P. E. Hartman, E. H. Beutner, and R. A. McLean. 1951a. Further evidence for the existence of mitochondria in bacteria. Journal of Bacteriology 62:729-739.
- Mudd, Stuart, Kenji Takeya, and Howard J. Henderson. 1956. Electron-scattering granules and reducing sites in mycobacteria. Journal of Bacteriology 72:767-783.
- Mudd, S., L. C. Winterscheid, E. D. Delama, and H. J. Henderson. 1951b. Evidence suggesting that the granules of mycobacteria are mitochondria. Journal of Bacteriology 62:459-475.

- Mudd, S., A. Yoshida, and M. Koide. 1958. Polyphosphate as accumulator of phosphorus and energy. Journal of Bacteriology 75:224-235.
- Munk, N., and H. Rosenberg. 1969. On the deposition and utilization of inorganic pyrophosphate in Tetrahymena pyriformis. Biochimica et Biophysica Acta 177:629-640.
- Murphy, J., and J. P. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. Analytica Chimica Acta 27:31-36.
- Ohle, W. 1953. Phosphor als Initialfaktor der Gewassereutrophierung. Vom Wasser 20:11-23.
- Overbeck, J. 1961a. The phosphatases of Scenedesmus quadricauda and their ecological importance. Verh. int. Vertin theor. angew. Limnol. 14:226-231.
- Overbeck, J. 1961b. Untersuchungen zum phosphatauchhalt von grunalgen - II. Die verwertung von pyrophosphat und organisch gebundenen phosphaten und ihre beziehung zu den phosphatesen von Scenedesmus quadricauda (Turp.) Breb. Aech. Hydrobiologie 58:281-308.
- Overbeck, J. 1962. Untersuchungen zum phosphat Naushalt von Grunalgen. III. Das Verhalten der zelefraktionen von Scenedesmus quadricauda (Turp.) Bre'b in Tagescyclus unter Verschiedenen Belichtungsbedingungen und bei Verschiedenen phosphatverbindungen, Archives fur Mikrobiologie 41:11-26.
- Pankratz, H. S., and C. C. Bowen. 1963. Cytology of blue-green algae. I. The cells of Symploca muscorum. American Journal of Botany 50:387-399.

- Pearsall, W. H. 1932. Phytoplankton in the English lakes. II. The composition of phytoplankton in relation to dissolved substances. Journal of Ecology 30:241-262.
- Pesch, K. L. 1924. Zentr. Bakteriolog. Parasitenk. Abt. I Orig. 92:208.
- Pilson, Michael E. Q., and Susan B. Betzer, 1973. Phosphorus flux across a coral reef. Ecology 54:581-588.
- Pitcairn, Carole E. R., and H. A. Hawkes. 1973. The role of phosphorus in the growth of Cladophora. Water Research 7:159-171.
- Powers, C. F., D. W. Schults, K. W. Malueg, Robert M. Brice, and Marcus D. Schuldt. 1972. Algal Responses to Nutrient Additions in Natural Waters. II. Field Experiments. In: Nutrients and Eutrophication: The Limiting Nutrient Controversy. G. E. Likens, ed. American Society of Limnology and Oceanography, Special Symposia, Volume I. pp 141-156.
- Reichardt, Wolfgang. 1971. Catalytic mobilization of phosphate in lake water and by Cyanophyta. Hydrobiologia 38:377-394.
- Reynolds, E. S. 1963. The use of lead citrate at high pH as an electron opaque stain in electron microscopy. Journal of Cell Biology 17:208-212.
- Rigler, F. H. 1956. A tracer study of the phosphorus cycle in lake water. Ecology 37:550-562.
- Rodhe, W. 1948. Environmental requirements of freshwater plankton algae. Symbol. Bot. Upsalienses 10:1-149.
- Rodhe, W. 1969. Crystallization of eutrophication concepts in northern Europe, p. 50-64. In: Eutrophication: Causes, consequences, and correctives. Nat. Acad. Sci./Nat. Res. Council, Publ. 1700.

- Rosenberg, H. 1966. The isolation and identification of "volutin" granules from Tetrahymena. Experimental Cell Research 41:397-410.
- Sakaguchi, Osamu, Katushi Yokota, and Tomoyūki Koshi. 1972. Acid and alkaline phosphatases of Vibrio parahaemolyticus. Japanese Journal of Microbiology 16:351-358.
- Sall, Theodore, Stuart Mudd, and John C. Davis. 1956. Factors conditioning the accumulation and disappearance of metaphosphate in cells of Corynebacterium diphtheriae. Archives of Biochemistry and Biophysics 60:130-146.
- Sall, Theodore, Stuart Mudd, and Atsushi Takagi. 1958. Polyphosphate accumulation and utilization as related to synchronized cell division of Corynebacterium diphtheriae. Journal of Bacteriology 76:640-645.
- Sauer, H. W., E. M. Goodman, K. L. Babcock, and H. P. Rusch. 1969. Polyphosphate in the life cycle of Physarum polycephalum and its relation to RNA synthesis. Biochimica et Biophysica Acta 195:401-409.
- Sawyer, C. N. 1947. Fertilization of lakes by agricultural and urban drainage. Journal of the New England Water Works Association 61:109-127.
- Sawyer, C. N. 1952. Some new aspects of phosphates in relation to lake fertilization. Sewage and Industrial Wastes 24:768-776.
- Schelske, Claire L., and Eugene F. Stoermer. 1972. Phosphorus, Silica, and Eutrophication of Lake Michigan. In: Nutrients and Eutrophication: The Limiting Nutrient Controversy. G. E.

- Likens, ed. American Society of Limnology and Oceanography, Special Symposia, Volume 1. pp 157-171.
- Schmidt, G., and S. J. Thannhauser. 1945. A method for the determination of desoxyribose nucleic acid, ribonucleic acid and phosphorproteins in animal tissues. Journal of Biological Chemistry 161:83-89.
- Schmidt, G., L. Hecht, and S. J. Thannhauser. 1946. The enzymatic formation and the accumulation of large amounts of a metaphosphate in bakers' yeast under certain conditions. Journal of Biological Chemistry 166:775-776.
- Schmidt, Gerhard, Liselotte Hecht, and S. J. Thannhauser. 1949. The effect of potassium ions on the absorption of orthophosphate and the formation of metaphosphate by bakers' yeast. Journal of Biological Chemistry 178:773-742.
- Schmidt, Gerhard, Knikor Seraidarian, Lowell M. Greenbaum, Mary D. Hickey, and S. J. Thannhauser. 1956. The effects of certain nutritional conditions on the formation of purines and of ribonucleic acid in Baker's yeast. Biochimica et Biophysica Acta 20:135-149.
- Schumacher, G. J., and L. A. Whitford. 1965. Respiration and ³²P uptake in various species of freshwater algae as affected by a current. Journal of Phycology 1:78-80.
- Shannon, James E., and G. Fred Lee. 1966. Hydrolysis of condensed phosphates in natural waters. Air and Water Pollution International Journal 10:735-756.

- Shapiro, Joseph. 1967. Induced Rapid Release and Uptake of Phosphate by Microorganisms. Science 155:1269-1271.
- Shapiro, Joseph. 1968. Studies on the natural factors affecting phosphate absorption and its utilization by algae. Water Research 2:21-23.
- Shapiro, Joseph. 1970. A statement on phosphorus. Journal of the Water Pollution Control Federation 42:772-775.
- Sicko, Linda M. 1972. Structural variations of polyphosphate bodies in blue-green algae. 30th Annual Proceedings of the Electron Microscopy Society of America, pp 218-219.
- Simonis, W., and W. Urbach. 1963. Uber eine Wirkung von Natriumionen auf die Phosphataufnahme und die lichtabhängige Phosphorylierung von Ankistrodesmus braunii. Archives fur Mikrobiologie 46:265-286.
- Solt, G. S. 1973. Phosphorus in industrial water. Water Research 7:69-76.
- Smith, I. W., J. F. Wilkinson, and J. P. Duguid. 1954. Volutin production in Aerobacter aerogenes due to nutrient imbalance. Journal of Bacteriology 68:450-463.
- Sonzogni, William C., and G. Fred Lee. 1972. Effect of Diversion of Domestic Waste Waters on Phosphorus Content and Eutrophication of the Madison Lakes. Report to the 73rd National Meeting, American Institute of Chemical Engineers, Minneapolis, Minnesota, August 1972.
- Spitznagel, J. K., and D. G. Sharp. 1959. Magnesium and sulfate ions as determinants in the growth and reproduction of Mycobacterium

- bovis. Journal of Bacteriology 78:453-562.
- Starr, R. C. 1964. The culture of algae at Indiana University.
American Journal of Botany 51:1010-1044.
- Stempak, J. F., and R. T. Ward. 1964. An improved staining method for
electron microscopy. Journal of Cell Biology 22:697-701.
- Stewart, W. D. P., and Gail Alexander. 1971. Phosphorus availability
and nitrogenase activity in aquatic blue-green algae.
Freshwater Biology 1:389-404.
- Suzuki, Hiroyasu, Takichi Kaneko, and Yonosuka Ikeda. 1972. Properties
of polyphosphate kinase prepared from Mycobacterium smegmatis.
Biochimica et Biophysica Acta 268:381-390.
- Sylvester, R. O. 1961. Nutrient content of drainage waters from
forested, urban and agricultural areas. In: Algae and
Metropolitan Wastes. pp 80-87. R. A. Taft Sanitary
Engineering Center, Cincinnati, Ohio. 162 pp.
- Talpasayi, E. R. S. 1962. Transients in radio-active phosphate
incorporation under nitrogen and argon by Anabaena cylindrica.
Plant and Cell Physiology 3:189-191.
- Talpasayi, E. R. S. 1963. Polyphosphate containing particles of
blue-green algae. Cytologia 28:76-80.
- Taylor, Alan W. 1967. Phosphorus and water pollution. Journal of
Soil and Water Conservation 11-12:228-231.
- Terry, K. R., and A. B. Hooper. 1970. Polyphosphate and orthophosphate
content of Nitrosomonas europaea as a function of growth.
Journal of Bacteriology 103:199-206.

- Thilo, E. 1962. Condensed phosphates and arsenates. Advances in Inorganic Chemistry and Radiochemistry 4:1-77.
- Thomas, E. A. 1953. Zur Bekämpfung der see - Eutrophierung: Empirische und experimentelle Untersuchungen zur Kenntnis der Minimumstoffe in 46 seen der schweiz und agrenzender gebiete. Monatsbull. Schweiz. Verein. Gas - Wasserfachm., No. 2-3. 15 p.
- Torriani, Annamaria. 1960. Influence of inorganic phosphate in the formation of phosphatases by Escherichia coli. Biochimica et Biophysica Acta 38:460-469.
- Ullrich, Wolfram R. 1972. Der Einfluß von CO₂ und pH auf die ³²P - Markierung von polyphosphaten und organischen phosphaten bei Ankistrodesmus braunii im Licht. Planta 102:37-54.
- Ullrich-Eberius, C. I. 1973. Die pH - Abhängigkeit der aufnahme von H₂PO₄⁻, SO₄⁼, Na⁺ und K⁺ und ihre gegenseitige Beeinflussung bei Ankistrodesmus braunii. Planta 109:161-176.
- Ullrich-Eberius, C. I., and W. Simonis. 1970. Der Einflup von Natrium - und kaliumionen auf die phosphataufnahme bei Ankistrodesmus braunii. Planta 93:214-226.
- Voelz, Herbert, Ursula Voelz, and Raoul O. Ortigoza. 1966. The "polyphosphate overplus" phenomenon in Myxococcus xanthus and its influence on the architecture of the cell. Archives fur Mikrobiologie 53:371-388.
- Wang, Dalton, and Dino Mancini. 1966. Studies on ribonucleic acid - polyphosphate in plants. Biochimica et Biophysica Acta 129:231-239.

- Weaver, P. J. 1969. Phosphates in surface waters and detergents. Journal of the Water Pollution Control Federation 41:1647-1653.
- Weibel, S. R., R. B. Weidner, J. M. Cohen, and A. G. Christianson. 1966. Pesticides and other contaminants in rainfall and runoff. Journal of the American Water Works Association 58:1075-1084.
- Weimberg, Ralph. 1970. Effect of potassium chloride on the uptake and storage of phosphate by Saccharomyces mellis. Journal of Bacteriology 103:37-48.
- Weimberg, Ralph, and William L. Orton. 1965. Synthesis and breakdown of the polyphosphate fraction and acid phosphomonoesterase of Saccharomyces mellis and their locations in the cell. Journal of Bacteriology 80:740-747.
- Whitton, B. A. 1967. Phosphate accumulation by colonies of Nostoc. Plant and Cell Physiology 8:293-296.
- Wiame, J. M. 1947a. The metachromatic reaction of hexametaphosphate. Journal of the American Chemical Society 69:3146-3147.
- Wiame, J. M. 1947b. Etude d'une substance polyphosphoree, basophile et metachromatique chez les levures. Biochimica et Biophysica Acta 1:234-255.
- Wiame, J. M. 1949. The occurrence and physiological behavior of two metaphosphate fractions in yeast. Journal of Biological Chemistry 178:919-929.
- Widra, Abe. 1959. Metachromatic granules of microorganisms. Journal of Bacteriology 78:664-670.

- Widra, A., and S. Wilburn. 1959. Electron microscopy of volutin granules in Aerobacter aerogenes. Giorn. Microbiology 7:81-84.
- Wilkinson, J. F., and J. P. Duguid. 1960. The influence of cultural conditions on bacterial cytology. International Review of Cytology 9:1-76.
- Winder, Frank, and Joan M. Denny. 1955. Utilization of Metaphosphate for phosphorylation by cell-free extracts of Mycobacterium smegmatis. Nature 175:636.
- Winder, F. G., and Joan M. Denny. 1956. Phosphorus metabolism of Mycobacteria: Determination of phosphorus compounds in some Mycobacteria. Journal of General Microbiology 15:1-18.
- Winder, F. G., and J. M. Denny. 1957. The metabolism of inorganic polyphosphate in Mycobacteria. Journal of General Microbiology 17:573-585.
- Yall, Irving, William H. Boughton, Richard C. Knudsen, and Norval A. Sinclair. 1970. Biological uptake of phosphorus by activated sludge. Applied Microbiology 20:145-150.
- Yoshida, A. 1955. Metaphosphate. II. Heat of hydrolysis of metaphosphate extracted from yeast cells. Journal of Biochemistry (Tokyo) 42:163-168.
- Zehnder, A., and P. R. Gorham. 1960. Factors influencing the growth of Microcystis aeruginosa Kutz. Emend Elenkin. Canadian Journal of Microbiology 6:645-660.